

ON THE CUSPIDAL SUPPORT OF DISCRETE SERIES FOR P-ADIC QUASISPLIT $Sp(N)$ AND $SO(N)$

BIN XU

ABSTRACT. Zelevinsky's classification theory of discrete series of p -adic general linear groups has been well known. Mœglin and Tadić gave the same kind of theory for p -adic classical groups, which is more complicated due to the occurrence of nontrivial structure of L-packets. Nonetheless, their work is independent of the endoscopic classification theory of Arthur (also Mok in the unitary case), which concerns the structure of L-packets in these cases. So our goal in this paper is to make more explicit the connection between these two very different types of theories. To do so, we reprove the results of Mœglin and Tadić in the case of quasisplit symplectic groups and orthogonal groups by using Arthur's theory.

1. INTRODUCTION

Let F be a p -adic field and G be a quasisplit connected reductive group over F . We consider pairs (M, π_{cusp}) for G , where M is a Levi subgroup of G and π_{cusp} is an irreducible supercuspidal representation of $M(F)$. Such pairs carry an action of $G(F)$ by conjugation, i.e.,

$$(M, \pi_{\text{cusp}})^g = (M^g, \pi_{\text{cusp}}^g),$$

where $M^g = g^{-1}Mg$, and $\pi_{\text{cusp}}^g(m) = \pi_{\text{cusp}}(gmg^{-1})$ for $m \in M(F)^g$. For any pair (M, π_{cusp}) , let $P = MN$ be any parabolic subgroup containing M , and we have the normalized parabolic induction $\text{Ind}_P^G(\pi_{\text{cusp}} \otimes 1_N)$. For simplicity we always abbreviate this to $\text{Ind}_P^G(\pi_{\text{cusp}})$, and we have the following facts about the parabolic induction.

- (1) $\text{Ind}_P^G(\pi_{\text{cusp}})$ is a smooth representation of finite length, i.e., the semi-simplification $s.s.\text{Ind}_P^G(\pi_{\text{cusp}})$ of $\text{Ind}_P^G(\pi_{\text{cusp}})$ is a direct sum of finitely many irreducible smooth representations.
- (2) For $g \in G(F)$, $s.s.\text{Ind}_P^G(\pi_{\text{cusp}}) = s.s.\text{Ind}_{P'}^G(\pi_{\text{cusp}}^g)$ for any parabolic subgroup P' containing M^g .

It is a theorem of Bernstein and Zelevinsky [BZ77] that all irreducible smooth representations of $G(F)$ can be constructed by parabolic induction from supercuspidal representations.

Theorem 1.1 (Bernstein-Zelevinsky). *For any irreducible smooth representation π of $G(F)$, there exists a unique pair (M, π_{cusp}) up to conjugation by $G(F)$ such that*

$$\pi \subseteq s.s.\text{Ind}_P^G(\pi_{\text{cusp}}).$$

Moreover, one can always find P' containing M such that

$$\pi \hookrightarrow \text{Ind}_{P'}^G(\pi_{\text{cusp}})$$

as a subrepresentation.

Remark 1.2. The $G(F)$ -conjugacy class of pairs (M, π_{cusp}) in this theorem is called the cuspidal support of π . For our later purposes, we would like to fix a Borel subgroup B of G together with a maximal torus $T \subseteq B$, and we have the standard parabolic subgroups $P = MN$, i.e., $P \supseteq B, M \supseteq T$. Then this theorem implies that for any irreducible smooth representation π of G , one can always find a standard parabolic subgroup $P = MN$ with a supercuspidal representation π_{cusp} of M such that $\pi \hookrightarrow \text{Ind}_P^G(\pi_{\text{cusp}})$ as a subrepresentation.

Based on this theorem, it is natural to ask the following questions.

2010 *Mathematics Subject Classification.* 22E50 (primary); 11F70 (secondary).

Key words and phrases. symplectic and orthogonal group, discrete series, cuspidal support, endoscopy.

Question 1.3. *How to determine the cuspidal support of any irreducible smooth representation of $G(F)$?*

Question 1.4. *How can one classify the irreducible **unitary** representations of $G(F)$ in terms of their cuspidal supports?*

Question 1.5. *How can one classify the irreducible **discrete series** representations of $G(F)$ in terms of their cuspidal supports?*

Question 1.3 is properly the most difficult one, and we are not able to say much about it here. Question 1.4 is often referred to as the **unitary dual problem**, and it has been solved for $GL(n)$ [Tad86]. For the classical groups, Tadić and Muić have done many works (see [Tad09], [MT11]), and again we will not say anything about it here. Our main interest is in Question 1.5, and it has the most complete theories for both $GL(n)$ (see [Zel80]) and classical groups (see [Mœg02], [MT02]). Our goal is to present the results for the quasisplit symplectic groups and special orthogonal groups. To be more precise about what we want to show, we need to introduce some notations.

If $G = GL(n)$, let us take B to be the group of upper-triangular matrices and T to be the group of diagonal matrices, then the standard Levi subgroup M can be identified with $GL(n_1) \times \cdots \times GL(n_r)$ through

$$\begin{pmatrix} GL(n_1) & & \\ & \ddots & \\ & & GL(n_r) \end{pmatrix}$$

$$(g_1, \dots, g_r) \longrightarrow \text{diag}\{g_1, \dots, g_r\}$$

with respect to any partition of $n = n_1 + \cdots + n_r$. So an irreducible supercuspidal representation π_{cusp} of $M(F)$ can be written as $\pi_{cusp} = \pi_1 \otimes \cdots \otimes \pi_r$ where π_i is an irreducible supercuspidal representation of $GL(n_i, F)$ for $1 \leq i \leq r$. For simplicity, we denote the normalized parabolic induction $\text{Ind}_P^G(\pi_{cusp})$ by $\pi_1 \times \cdots \times \pi_r$. An irreducible supercuspidal representation π of $GL(n, F)$ can always be written in a unique way as $\rho||^x := \rho \otimes |\det(\cdot)|^x$ for an irreducible unitary supercuspidal representation ρ and a real number x . To fix notations, we will always denote by ρ an irreducible unitary supercuspidal representation of $GL(d_\rho, F)$. Now for a finite length arithmetic progression of real numbers of common length 1 or -1

$$x, \dots, y$$

and an irreducible unitary supercuspidal representation ρ of $GL(d_\rho)$, it is a general fact that

$$\rho||^x \times \cdots \times \rho||^y$$

has a unique irreducible subrepresentation, denoted by $\langle \rho; x, \dots, y \rangle$ or $\langle x, \dots, y \rangle$. If $x \geq y$, it is called a Steinberg representation; if $x < y$, it is called a Speh representation. Such sequence of ordered numbers is called a **segment** (cf. Appendix B). In particular, when $x = -y > 0$, we can let $a = 2x + 1 \in \mathbb{Z}$ and write

$$St(\rho, a) := \langle \frac{a-1}{2}, \dots, -\frac{a-1}{2} \rangle,$$

which is an irreducible smooth representation of $GL(ad_\rho, F)$. In fact it is a discrete series representation by Zelevinsky's classification theorem.

Theorem 1.6 (Zelevinsky [Zel80]). *All irreducible discrete series representations of $GL(n, F)$ can be obtained in a unique way as $St(\rho, a)$ for certain irreducible unitary subcuspidal representation of $GL(d_\rho)$ and integer a so that $n = ad_\rho$.*

If $G = Sp(2n)$, let us define it with respect to

$$\begin{pmatrix} 0 & -J_n \\ J_n & 0 \end{pmatrix},$$

where

$$J_n = \begin{pmatrix} & & 1 \\ & \ddots & \\ 1 & & \end{pmatrix}.$$

We take B to be subgroup of upper-triangular matrices in G and T to be subgroup of diagonal matrices in G , then the standard Levi subgroup M can be identified with $GL(n_1) \times \cdots \times GL(n_r) \times G_-$ for any partition $n = n_1 + \cdots + n_r + n_-$ and $G_- = Sp(2n_-)$ as follows

$$(1.1) \quad \begin{pmatrix} GL(n_1) & & & & & 0 \\ & \ddots & & & & \\ & & GL(n_r) & & & \\ & & & G_- & & \\ & & & & GL(n_r) & \\ & & & & & \ddots \\ 0 & & & & & & GL(n_1) \end{pmatrix} (g_1, \dots, g_r, g) \longrightarrow \text{diag}\{g_1, \dots, g_r, g, {}_t g_r^{-1}, \dots, {}_t g_1^{-1}\},$$

where ${}_t g_i = J_{n_i} {}^t g_i J_{n_i}^{-1}$ for $1 \leq i \leq r$. Note n_- can be 0, in which case we simply write $Sp(0) = 1$. So an irreducible supercuspidal representation π_{cusp} of $M(F)$ can be written as $\pi_{\text{cusp}} = \pi_1 \otimes \cdots \otimes \pi_r \otimes \sigma$ where π_i is an irreducible supercuspidal representation of $GL(n_i, F)$ for $1 \leq i \leq r$ and σ is an irreducible supercuspidal representation of $G_-(F)$. For simplicity, we denote $\text{Ind}_P^G(\pi_{\text{cusp}})$ by $\pi_1 \times \cdots \times \pi_r \rtimes \sigma$. Note that σ is always unitary. The discussion here can be easily extended to special orthogonal groups.

If $G = SO(N)$ split, we can define it with respect to J_N . When N is odd, the situation is exactly the same as the symplectic case. When $N = 2n$, there are two distinctions. First, the standard Levi subgroups given through the embedding (1.1) do not exhaust all standard Levi subgroups of $SO(2n)$. To get all of them, we need to take the θ_0 -conjugate of M given in (1.1), where

$$(1.2) \quad \theta_0 = \begin{pmatrix} 1 & & & & \\ & \ddots & & & \\ & & 1 & & \\ & & & 1 & \\ & & & & \ddots \\ & & & & & 1 \end{pmatrix}.$$

Note $M^{\theta_0} \neq M$ only when $n_- = 0$ and $n_r > 1$. In order to distinguish the θ_0 -conjugate standard Levi subgroups of $SO(2n)$, we will only identify those Levi subgroups M in (1.1) with $GL(n_1) \times \cdots \times GL(n_r) \times G_-$, and we denote the other simply by M^{θ_0} . Second, if the partition $n = n_1 + \cdots + n_r + n_-$ satisfies $n_r = 1$ and $n_- = 0$, then we can rewrite it as $n = n_1 + \cdots + n_{r-1} + n'_-$ with $n'_- = 1$, and the corresponding Levi subgroup is the same. This is because $GL(1) \cong SO(2)$.

In this paper, we will also consider $G = SO(2n, \eta)$, which is the outer form of the split $SO(2n)$ with respect to a quadratic extension E/F and conjugation by θ_0 . Here η is the associated quadratic character of E/F by the local class field theory. Then the standard Levi subgroups of $SO(2n, \eta)$ will be the outer form of those θ_0 -stable standard Levi subgroups of $SO(2n)$. In particular, they can be identified with $GL(n_1) \times \cdots \times GL(n_r) \times SO(n_-, \eta)$ with $n_- \neq 0$.

Finally for any irreducible discrete series representation π of a symplectic group or special orthogonal group $G(F)$, our goal is to find unitary supercuspidal representations ρ_i of $GL(d_{\rho_i}, F)$ for $1 \leq i \leq r$ together with real numbers $x_1, \dots, x_r$, and a supercuspidal representation σ of $G_-(F)$ which is of the same type as G , such that

$$\pi \text{ or } \pi^{\theta_0} \hookrightarrow \rho_1 ||^{x_1} \times \cdots \times \rho_r ||^{x_r} \rtimes \sigma$$

as a subrepresentation.

The approach that we are going to take will highly rely on Arthur's endoscopic classification theory for symplectic and orthogonal groups [Art13], especially the structure of tempered Arthur packets (or L-packets). It is different from the original approaches of Mœglin and Tadić (see [Mœg02], [MT02]), where although possibly motivated by the structure of L-packets, they do not need to use it in their arguments. There are two reasons for us to adopt the new approach. One is there are certain reducibility assumptions (see Proposition 3.2) taken in the works of Mœglin and Tadić that could be removed under Arthur's work (as suggested by them), so it would be very natural to start with Arthur's theory at the first place. The other reason is the endoscopic theory is "hidden" in their works, but we want to see how it could play a role in this kind of classification theory, to be more precise, the interplay of endoscopy theory with the theory of Jacquet modules (see Section 3).

Acknowledgements: This work came out of the author's talks in the automorphic forms seminar at Fields Institute in 2014. The author wants to thank the organizer Chung Pang Mok for initially asking him to give those talks, otherwise this paper would not appear. The author also wants to thank all the participants in the seminar for their interests and supports. While writing up the current version of this paper, the author has enjoyed the hospitality of Institute for Advanced Study (IAS). He also gratefully acknowledges the support from University of Calgary and the Pacific Institute for the Mathematical Sciences (PIMS), when this work took its final form. At last, he wants to thank the referee for many helpful comments and suggestions. During his time at IAS, he was supported by the NSF under Grant No. DMS-1128155 and DMS-1252158.

2. TEMPERED ARTHUR PACKET

Let F be a p -adic field and G be a quasisplit symplectic group or special orthogonal group. We define the local Langlands group as $L_F = W_F \times SL(2, \mathbb{C})$, where W_F is the usual Weil group. We write $\Gamma_F = \Gamma_{\bar{F}/F}$ for the absolute Galois group over F . Let $\hat{G}$ be the complex dual group of G , and ${}^L G$ be the Langlands dual group of G . A tempered (or generic) Arthur parameter of G is a $\hat{G}$ -conjugacy class of admissible homomorphisms $\phi : L_F \rightarrow {}^L G$, such that $\phi|_{W_F}$ is bounded. We denote by $\Phi_{bdd}(G)$ the set of tempered Arthur parameters. Here we can simplify the Langlands dual groups as in the following table:

G	${}^L G$
$Sp(2n)$	$SO(2n+1, \mathbb{C})$
$SO(2n+1)$	$Sp(2n, \mathbb{C})$
$SO(2n, \eta)$	$SO(2n, \mathbb{C}) \rtimes \Gamma_{E/F}$

In the last case, η is a quadratic character associated with a quadratic extension E/F and $\Gamma_{E/F}$ is the associated Galois group. If we define $O(2n, \mathbb{C})$ with respect to J_{2n} , we can fix an isomorphism $SO(2n, \mathbb{C}) \rtimes \Gamma_{E/F} \cong O(2n, \mathbb{C})$ by sending the nontrivial element of $\Gamma_{E/F}$ to the permutation matrix (1.2). So in either of these cases, there is a natural embedding ξ_N of ${}^L G$ into $GL(N, \mathbb{C})$ up to $GL(N, \mathbb{C})$ -conjugacy, where $N = 2n+1$ if $G = Sp(2n)$ or $N = 2n$ otherwise. Under such an embedding, we can view the parameter ϕ as an equivalence class of N -dimensional self-dual representations of L_F , i.e., $\phi^\vee = \phi$. Let π_ϕ be the self-dual representation of $GL(N, F)$ associated with ϕ under the local Langlands correspondence (cf. [HT01], [Hen00], and [Sch13]). If we decompose ϕ into equivalence classes of irreducible subrepresentations, we get

$$(2.1) \quad \phi = \bigoplus_{i=1}^q l_i \phi_i,$$

where ϕ_i is an equivalence class of irreducible representations of L_F and l_i is the multiplicity. Since L_F is a product of W_F and $SL(2, \mathbb{C})$, we can further decompose ϕ_i as an tensor product

$$\phi_i = \phi_{cusp, i} \otimes \nu_{a_i},$$

where $\phi_{\text{cusp},i}$ is an equivalence class of irreducible unitary representations of W_F and ν_{a_i} is the $(a_i - 1)$ -th symmetric power representation of $SL(2, \mathbb{C})$. Now we have obtained all the combinatorial data needed from ϕ in the paper. To put it in a nice way, we first identify the set of equivalence classes of irreducible unitary supercuspidal representations of $GL(d, F)$ with equivalence classes of d -dimensional irreducible unitary representations of W_F through the local Langlands correspondence for $GL(d)$, and denote by ρ_i the corresponding representation for $\phi_{\text{cusp},i}$. Also notice the representation ν_{a_i} is completely determined by its dimension. So altogether we can simply write $\phi_i = \rho_i \otimes [a_i]$ formally. After this discussion we can define the multi-set of Jordan blocks for ϕ as follows,

$$Jord(\phi) := \{(\rho_i, a_i) \text{ with multiplicity } l_i : 1 \leq i \leq q\},$$

and

$$Jord_\rho(\phi) := \{a_i \text{ with multiplicity } l_i : \rho = \rho_i\}.$$

To parametrize the discrete series representations, we need to introduce a subset $\Phi_2(G)$ of $\Phi_{bdd}(G)$. Define

$$\Phi_2(G) := \{\phi \in \Phi_{bdd}(G) : \phi = \bigoplus_{i=1}^q \phi_i, \phi_i^\vee = \phi_i\}.$$

It is clear that the defining condition for $\Phi_2(G)$ is equivalent to requiring $Jord(\phi)$ is multiplicity free and $Jord_\rho(\phi)$ is empty unless ρ is self-dual. Moreover, for certain parity reason (see Section 3) the integers in $Jord_\rho(\phi)$ must be all odd or all even when $\phi \in \Phi_2(G)$. Besides, there is another description of $\Phi_2(G)$. For $\phi \in \Phi_{bdd}(G)$, we fix a representative $\underline{\phi}$. Let us define

$$\begin{aligned} S_{\underline{\phi}} &= \text{Cent}(\text{Im } \underline{\phi}, \widehat{G}), \\ \bar{S}_{\underline{\phi}} &= S_{\underline{\phi}} / Z(\widehat{G})^{\Gamma_F}, \\ \mathcal{S}_{\underline{\phi}} &= \bar{S}_{\underline{\phi}} / \bar{S}_{\underline{\phi}}^0 = S_{\underline{\phi}} / S_{\underline{\phi}}^0 Z(\widehat{G})^{\Gamma_F}. \end{aligned}$$

Then we have the following fact.

Lemma 2.1. *For $\phi \in \Phi_{bdd}(G)$, $\phi \in \Phi_2(G)$ if and only if $\bar{S}_{\underline{\phi}}$ is finite.*

This lemma can be shown by computing the group $S_{\underline{\phi}}$ explicitly (see [Art13, Section 1.4]). In particular, one can show $\mathcal{S}_{\underline{\phi}}$ is abelian.

To state Arthur's classification theory of tempered representations of quasisplit symplectic and orthogonal groups, we need to introduce some more notations. We will fix an outer automorphism θ_0 of G preserving an F -splitting. If G is symplectic or special odd orthogonal, we let $\theta_0 = id$. If G is special even orthogonal, we let θ_0 be induced from the conjugate action of the permutation matrix (1.2). Let $\widehat{\theta}_0$ be the dual automorphism of θ_0 . We write $\Sigma_0 = \langle \theta_0 \rangle$, $G^{\Sigma_0} = G \rtimes \Sigma_0$, and $\widehat{G}^{\Sigma_0} = \widehat{G} \rtimes \langle \widehat{\theta} \rangle$. Let ω_0 be the character of G^{Σ_0}/G , which is nontrivial when G is special even orthogonal. So in the special even orthogonal case, we can send θ_0 to the permutation matrix (1.2) to get an isomorphism between G^{Σ_0} and the full orthogonal group. If G has $\bar{F}$ -rank n , we write $G = G(n)$. Let $G(0) = G(0)^{\Sigma_0} = 1$. Also for the trivial representation of $G(0)$, we require formally $1^{\theta_0} \not\cong 1$ if $G(0) = SO(0)$, and $1^{\theta_0} \cong 1$ otherwise.

Let $\Pi_{\text{temp}}(G)$ (resp. $\Pi_2(G)$) be the set of equivalence classes of irreducible tempered representations (resp. discrete series representations) of $G(F)$. Note Σ_0 acts on these sets by conjugation, and we denote the set of Σ_0 -orbits in $\Pi_{\text{temp}}(G)$ (resp. $\Pi_2(G)$) by $\bar{\Pi}_{\text{temp}}(G)$ (resp. $\bar{\Pi}_2(G)$). Also note Σ_0 acts on $\Phi_{bdd}(G)$ (resp. $\Phi_2(G)$) through $\widehat{\theta}_0$, and we denote the corresponding set of Σ_0 -orbits by $\bar{\Phi}_{bdd}(G)$ (resp. $\bar{\Phi}_2(G)$). It is clear that for $\phi \in \Phi_{bdd}(G)$, $Jord(\phi)$ only depends on its image in $\bar{\Phi}_{bdd}(G)$. It is because of this reason, we will also denote the elements in $\bar{\Phi}_{bdd}(G)$ by ϕ . Moreover, through the natural embedding ξ_N , we can view $\bar{\Phi}_{bdd}(G)$ as a subset of equivalence classes of N -dimensional self-dual representations of L_F .

Theorem 2.2 (Arthur [Art13], Theorem 1.5.1). (1) *For $\phi \in \bar{\Phi}_{bdd}(G)$, one can associate a finite set $\bar{\Pi}_\phi$ of $\bar{\Pi}_{\text{temp}}(G)$, determined by π_ϕ through the theory of twisted endoscopy (see Section 4). And*

for a fixed Whittaker datum, there is a canonical bijection between $\bar{\Pi}_\phi$ and characters $\widehat{\mathcal{S}}_\phi$ of $\mathcal{S}_\phi$.

$$\bar{\Pi}_\phi \longrightarrow \widehat{\mathcal{S}}_\phi$$

$$[\pi] \longmapsto \langle \cdot, \pi \rangle_\phi$$

(2) *There are decompositions*

$$\begin{aligned} \bar{\Pi}_{temp}(G) &= \bigsqcup_{\phi \in \bar{\Phi}_{bdd}(G)} \bar{\Pi}_\phi, \\ \bar{\Pi}_2(G) &= \bigsqcup_{\phi \in \bar{\Phi}_2(G)} \bar{\Pi}_\phi. \end{aligned}$$

We will denote the characters of $\mathcal{S}_\phi$ by $\bar{\varepsilon}$, and denote the corresponding Σ_0 -orbit $[\pi]$ of irreducible representations by $\pi(\phi, \bar{\varepsilon})$. Let us define $\Pi_\phi^{\Sigma_0}$ to be set of irreducible representations of $G^{\Sigma_0}(F)$ whose restriction to $G(F)$ belong to $\bar{\Pi}_\phi$. We call an irreducible representation π^{Σ_0} of $G^{\Sigma_0}(F)$ is a discrete series if its restriction to $G(F)$ are discrete series representations. We also define $\mathcal{S}_\phi^{\Sigma_0}$, $\bar{\mathcal{S}}_\phi^{\Sigma_0}$ and $\widehat{\mathcal{S}}_\phi^{\Sigma_0}$ as before simply by taking $\widehat{G}^{\Sigma_0}$ in place of $\widehat{G}$. The following theorem asserts $\Pi_\phi^{\Sigma_0}$ can be parametrized by the characters of $\mathcal{S}_\phi^{\Sigma_0}$. It is a consequence of [Art13, Theorem 2.2.4].

Theorem 2.3 (Arthur). *Suppose $\phi \in \bar{\Phi}_{bdd}(G)$, for a fixed Σ_0 -stable Whittaker datum there is a canonical bijection between $\Pi_\phi^{\Sigma_0}$ and characters $\widehat{\mathcal{S}}_\phi^{\Sigma_0}$ of $\mathcal{S}_\phi^{\Sigma_0}$*

$$\Pi_\phi^{\Sigma_0} \longrightarrow \widehat{\mathcal{S}}_\phi^{\Sigma_0}$$

$$\pi^{\Sigma_0} \longmapsto \langle \cdot, \pi^{\Sigma_0} \rangle_\phi,$$

such that

$$\langle \cdot, \pi^{\Sigma_0} \rangle_\phi|_{\mathcal{S}_\phi} = \langle \cdot, \pi \rangle_\phi,$$

where $\pi \subseteq \pi^{\Sigma_0}|_G$.

We denote the characters of $\mathcal{S}_\phi^{\Sigma_0}$ by ε , and denote the corresponding representations by $\pi^{\Sigma_0}(\phi, \varepsilon)$. We also denote the image of ε in $\widehat{\mathcal{S}}_\phi$ by $\bar{\varepsilon}$. If ε_0 is the generator of the kernel of the projection $\widehat{\mathcal{S}}_\phi^{\Sigma_0} \rightarrow \widehat{\mathcal{S}}_\phi$, then this theorem implies

$$(2.2) \quad \pi^{\Sigma_0}(\phi, \varepsilon \varepsilon_0) \cong \pi^{\Sigma_0}(\phi, \varepsilon) \otimes \omega_0.$$

Therefore, if G is special even orthogonal and $\mathcal{S}_\phi^{\Sigma_0} \neq \mathcal{S}_\phi$, then $\pi(\phi, \bar{\varepsilon})$ is a representation of $G(F)$ satisfying $\pi(\phi, \bar{\varepsilon})^{\theta_0} \cong \pi(\phi, \bar{\varepsilon})$. The converse is also true, i.e., if G is special even orthogonal and $\mathcal{S}_\phi^{\Sigma_0} = \mathcal{S}_\phi$, then $\pi^{\theta_0} \not\cong \pi$ for $\pi(\phi, \bar{\varepsilon}) = [\pi]$. In the rest of this paper, we will always fix a Σ_0 -stable Whittaker datum of G .

3. PARAMETERS OF SUPERCUSPIDAL REPRESENTATIONS

We keep the notations from the previous section.

Proposition 3.1. *Suppose π is a supercuspidal representation of $G(F)$, and $[\pi] \in \bar{\Pi}_\phi$ for some $\phi \in \bar{\Phi}_2(G)$. Then if $(\rho, a) \in \text{Jord}(\phi)$, one must have $(\rho, a-2) \in \text{Jord}(\phi)$ as long as $a-2 > 0$.*

Proof. Let ρ be a unitary irreducible supercuspidal representation of $GL(d_\rho, F)$. We can view $GL(d_\rho) \times G$ as the Levi component M_+ of a standard maximal parabolic subgroup P_+ of G_+ , where G and G_+ are of the same type. Let $\pi_{M_+} = \rho \otimes \pi$, and w is the unique non-trivial element in the relative Weyl group $W(M_+, G_+)$, which acts on $GL(d_\rho)$ as an outer automorphism. Let $\pi_{M_+, \lambda} = \rho ||^\lambda \otimes \pi$ for $\lambda \in \mathbb{C}$. It is a result of Arthur (see [Art13, Section 2.3]) that for certain choice of representative $\dot{w}$ of w , the standard intertwining operator between $\text{Ind}_{P_+}^{G_+}(\pi_{M_+, \lambda})$ and $\text{Ind}_{P_+}^{G_+}(\dot{w} \pi_{M_+, \lambda})$, i.e.,

$$J_{P_+}(\dot{w}, \pi_{M_+, \lambda})h(g) = \int_{N_{P_+} \cap wN_{P_+}w^{-1} \backslash N_{P_+}} h(\dot{w}^{-1}ng)dn, \quad h \in \text{Ind}_{P_+}^{G_+}(\pi_{M_+, \lambda}),$$

and the standard intertwining operator $J_{P_+}(\dot{w}^{-1}, \dot{w} \pi_{M_+, \lambda})$ between $\text{Ind}_{P_+}^{G_+}(\dot{w} \pi_{M_+, \lambda})$ and $\text{Ind}_{P_+}^{G_+}(\pi_{M_+, \lambda})$ can be normalized by meromorphic functions $r_{P_+}(w, \phi_{M_+, \lambda})$ and $r_{P_+}(w^{-1}, w \phi_{M_+, \lambda})$ respectively, i.e.,

$$\begin{aligned} R_{P_+}(\dot{w}, \pi_{M_+, \lambda}) &:= r_{P_+}(w, \phi_{M_+, \lambda})^{-1} J_{P_+}(\dot{w}, \pi_{M_+, \lambda}), \\ R_{P_+}(\dot{w}^{-1}, \dot{w} \pi_{M_+, \lambda}) &:= r_{P_+}(w^{-1}, w \phi_{M_+, \lambda})^{-1} J_{P_+}(\dot{w}^{-1}, \dot{w} \pi_{M_+, \lambda}), \end{aligned}$$

so that

$$(3.1) \quad R_{P_+}(\dot{w}^{-1}, \dot{w} \pi_{M_+, \lambda}) R_{P_+}(\dot{w}, \pi_{M_+, \lambda}) = Id.$$

Here $\phi_{M_+, \lambda}$ denotes the Langlands parameter for $\pi_{M_+, \lambda}$, and

$$r_{P_+}(w, \phi_{M_+, \lambda}) \sim \frac{L(\lambda, \rho \times \pi_\phi) L(2\lambda, \rho, R)}{L(1 + \lambda, \rho \times \pi_\phi) L(1 + 2\lambda, \rho, R)}$$

where R is either a symmetric square (S^2) or a skew-symmetric square ($\wedge^2$) representation of $GL(d_\rho, \mathbb{C})$ and “ $\sim$ ” means equal up to a nowhere vanishing holomorphic function of λ (that is given by the ϵ -factors here). Note $R = \wedge^2$ if G is $Sp(2n)$, $SO(2n, \eta)$ or $R = S^2$ if $G = SO(2n + 1)$. Similarly we have

$$r_{P_+}(w^{-1}, w \phi_{M_+, \lambda}) \sim \frac{L(-\lambda, \rho^\vee \times \pi_\phi) L(-2\lambda, \rho^\vee, R)}{L(1 - \lambda, \rho^\vee \times \pi_\phi) L(1 - 2\lambda, \rho^\vee, R)}.$$

Then we can rewrite (3.1) as

$$(3.2) \quad J_{P_+}(\dot{w}^{-1}, \dot{w} \pi_{M_+, \lambda}) J_{P_+}(\dot{w}, \pi_{M_+, \lambda}) \sim \frac{L(\lambda, \rho \times \pi_\phi) L(2\lambda, \rho, R) L(-\lambda, \rho^\vee \times \pi_\phi) L(-2\lambda, \rho^\vee, R)}{L(1 + \lambda, \rho \times \pi_\phi) L(1 + 2\lambda, \rho, R) L(1 - \lambda, \rho^\vee \times \pi_\phi) L(1 - 2\lambda, \rho^\vee, R)}.$$

Since π_{M_+} is supercuspidal, it follows from a theorem of Harish-Chandra ([Sil79, Theorem 5.4.2.1] and [Sha81, Lemma 2.2.5]) that both $J_{P_+}(\dot{w}, \pi_{M_+, \lambda})$ and $J_{P_+}(\dot{w}^{-1}, \dot{w} \pi_{M_+, \lambda})$ are holomorphic for $\text{Re } \lambda \neq 0$. So now we will assume $\lambda \in \mathbb{R}$ and $\lambda > 1/2$. Since $L(s, \rho \times \pi_\phi)$ does not have a pole (non-vanishing is clear) for real $s > 0$, and $L(s, \rho, R)$ does not have a pole (non-vanishing is clear) for real $s \neq 0$, we have

$$(3.3) \quad J_{P_+}(\dot{w}^{-1}, \dot{w} \pi_{M_+, \lambda}) J_{P_+}(\dot{w}, \pi_{M_+, \lambda}) \sim \frac{L(-\lambda, \rho^\vee \times \pi_\phi)}{L(1 - \lambda, \rho^\vee \times \pi_\phi)} \quad (\lambda > 1/2).$$

Finally, we learn from the definition of $L(s, \rho^\vee \times \pi_\phi)$ that it has a pole at $s = -(a-1)/2$ if and only if $\rho \cong \rho^\vee$ and $(\rho, a) \in \text{Jord}(\phi)$ (see Appendix A). So if we know $(\rho, a) \in \text{Jord}(\phi)$ for $a > 2$, then $L(-\lambda, \rho^\vee \times \pi_\phi)$ has a pole at $\lambda = (a-1)/2 > 1/2$. By the holomorphy of standard intertwining operators on the left hand side of (3.3), $L(1 - \lambda, \rho^\vee \times \pi_\phi)$ must also have a pole at $\lambda = (a-1)/2$, i.e., $1 - \lambda = 1 - (a-1)/2 = -(a-3)/2$ is a pole of $L(s, \rho^\vee \times \pi_\phi)$. So $(\rho, a-2) \in \text{Jord}(\phi)$. $\square$

If ρ is a self-dual unitary irreducible supercuspidal representation of $GL(d_\rho, F)$, we know from Appendix A that

$$L(s, \rho \times \rho) = L(s, \rho, \wedge^2) L(s, \rho, S^2)$$

has a pole at $s = 0$. We call ρ is of **symplectic type** if $L(s, \rho, \wedge^2)$ has a pole at $s = 0$, and we call ρ is of **orthogonal type** if $L(s, \rho, S^2)$ has a pole at $s = 0$. Moreover, for any positive integer a , the pair (ρ, a) is called having orthogonal type if ρ is of orthogonal type and a is odd, or ρ is of symplectic

type and a is even. Otherwise (ρ, a) is called having symplectic type. Next we are going to prove a very important reducibility result, which is named “Basic Assumption” in [Mœg02], [MT02]. Those careful readers may notice there is a slight difference between our statement below and the original one. The reason is they consider the group G^{Σ_0} rather than G , nonetheless one can translate between these two statements without difficulty (see Corollary 9.1).

Proposition 3.2. *Suppose π is a supercuspidal representation of $G(F)$, and $[\pi] \in \bar{\Pi}_\phi$ for some $\phi \in \bar{\Phi}_2(G)$. Then for any unitary irreducible supercuspidal representation ρ of $GL(d_\rho, F)$ and real number a_ρ , the parabolic induction*

$$\rho ||^{\pm(a_\rho+1)/2} \rtimes \pi$$

reduces if and only if ρ is self-dual and

$$(3.4) \quad a_\rho = \begin{cases} \max \text{Jord}_\rho(\phi), & \text{if } \text{Jord}_\rho(\phi) \neq \emptyset, \\ 0, & \text{if } \text{Jord}_\rho(\phi) = \emptyset, \rho \text{ is of opposite type to } \widehat{G}, \\ -1, & \text{otherwise, provided } d_\rho \text{ is even or } \pi \cong \pi^{\theta_0}. \end{cases}$$

Proof. We will follow the proof of the previous proposition and let $\lambda \in \mathbb{R}$. Suppose $\lambda > 0$. Then the image of $J_{P_+}(\dot{w}, \pi_{M_+, \lambda})$ is nonzero and irreducible by [BW00, Proposition 2.6]. It follows $\rho ||^\lambda \rtimes \pi$ is irreducible if and only if the kernel of $J_{P_+}(\dot{w}, \pi_{M_+, \lambda})$ is trivial. Since $\text{Ind}_{P_+}^{G_+}(\dot{w} \pi_{M_+, \lambda})$ and $\text{Ind}_{P_+}^{G_+}(\pi_{M_+, \lambda})$ have the same irreducible constituents (see [BZ77, Theorem 2.9]), the kernel of $J_{P_+}(\dot{w}, \pi_{M_+, \lambda})$ is trivial if and only if $J_{P_+}(\dot{w}, \pi_{M_+, \lambda})$ is an isomorphism. As we have seen previously, $J_{P_+}(\dot{w}, \pi_{M_+, \lambda})$ and $J_{P_+}(\dot{w}^{-1}, \dot{w} \pi_{M_+, \lambda})$ are holomorphic. In fact, they are also nonzero (see [Wal03, Section 4.1]). So $J_{P_+}(\dot{w}, \pi_{M_+, \lambda})$ is an isomorphism if and only if

$$J_{P_+}(\dot{w}^{-1}, \dot{w} \pi_{M_+, \lambda}) J_{P_+}(\dot{w}, \pi_{M_+, \lambda}) \neq 0.$$

As a consequence, $\rho ||^\lambda \rtimes \pi$ is reducible if and only if

$$J_{P_+}(\dot{w}^{-1}, \dot{w} \pi_{M_+, \lambda}) J_{P_+}(\dot{w}, \pi_{M_+, \lambda}) = 0.$$

Let us first assume $\lambda > 1/2$, then from (3.3) it is enough to see when

$$(3.5) \quad \frac{L(-\lambda, \rho^\vee \times \pi_\phi)}{L(1 - \lambda, \rho^\vee \times \pi_\phi)} = 0,$$

i.e., $L(1 - \lambda, \rho^\vee \times \pi_\phi)$ has a pole, but $L(-\lambda, \rho^\vee \times \pi_\phi)$ does not. From our discussion in the previous proof we know this can only happen when $\rho = \rho^\vee$ and $\lambda = (a_\rho + 1)/2$, where a_ρ is $\max \text{Jord}_\rho(\phi)$. Next we assume $0 < \lambda \leq 1/2$, it follows from (3.2) that

$$J_{P_+}(\dot{w}^{-1}, \dot{w} \pi_{M_+, \lambda}) J_{P_+}(\dot{w}, \pi_{M_+, \lambda}) \sim \frac{L(-\lambda, \rho^\vee \times \pi_\phi)}{L(1 - 2\lambda, \rho^\vee, R)}.$$

And the right hand side can be zero only when $L(1 - 2\lambda, \rho^\vee, R)$ has a pole, but $L(-\lambda, \rho^\vee \times \pi_\phi)$ does not. So necessarily $\rho = \rho^\vee$ and $\lambda = 1/2$. By our assumption on the representation R , we know $L(s, \rho, R)$ has a pole at $s = 0$ if and only if ρ is of opposite type to $\widehat{G}$. And the requirement that $L(s, \rho \times \pi_\phi)$ does not have a pole at $-1/2$ implies $\text{Jord}_\rho(\phi) = \emptyset$.

For $\lambda < 0$, one just needs to notice $s.s.(\rho ||^s \rtimes \pi) = s.s.(\rho^\vee ||^{-s} \rtimes \pi)^\theta$ for some $\theta \in \Sigma_0$, so one can apply the same argument to $\rho^\vee ||^{-\lambda} \rtimes \pi$.

Finally, we consider $\lambda = 0$, where our previous criterion does not work. However the reducibility of $\rho \rtimes \pi$ follows from the standard theory of representation theoretic R -groups. In Arthur’s theory these groups have been shown to be isomorphic to R -groups defined by parameters, which can be computed explicitly (see [Art13, Section 2.4 and Section 6.6]). So our reducibility condition in this case will follow from there. $\square$

Suppose π is an irreducible supercuspidal representation of $G(F)$ and $[\pi] \in \bar{\Pi}_\phi$ for some $\phi \in \bar{\Phi}_2(G)$. We know from Proposition 3.1 that $\text{Jord}(\phi)$ should be in a certain shape, and in view of Theorem 2.2 we would also like to know what kind of character $\bar{\varepsilon}$ of $\mathcal{S}_\phi$ will parametrize $[\pi]$. To give a description of such

characters, we have to first make an identification between $\widehat{\mathcal{S}}_{\underline{\phi}}^{\Sigma_0}$ with $\mathbb{Z}_2$ -valued functions over $Jord(\phi)$. To be more precise, let us assume

$$(3.6) \quad \underline{\phi} = \underline{\phi}_1 \oplus \cdots \oplus \underline{\phi}_r$$

where $\underline{\phi}_i$ are self-dual irreducible representations of dimension n_i . By Schur's Lemma,

$$\text{Cent}(\underline{\phi}, GL(N, \mathbb{C})) \cong \underbrace{\mathbb{C}^\times \times \cdots \times \mathbb{C}^\times}_r$$

where each $\mathbb{C}^\times$ acts on the corresponding representation space of $\underline{\phi}_i$. So

$$\text{Cent}(\underline{\phi}, \widehat{G}) \cong \{s = (s_i) \in \mathbb{Z}_2^r : \prod_i (s_i)^{n_i} = 1\}.$$

Note $\mathcal{S}_{\underline{\phi}}^{\Sigma_0} = \bar{\mathcal{S}}_{\underline{\phi}}^{\Sigma_0}$ in this case. Then $\mathcal{S}_{\underline{\phi}}^{\Sigma_0} \cong \mathbb{Z}_2^r / \langle (-1, \dots, -1) \rangle$. Since the right hand side does not depend on the choice of representative $\underline{\phi}$, we can denote it by $\mathcal{S}_{\phi}^{\Sigma_0}$. If G is special even orthogonal,

$$\mathcal{S}_{\underline{\phi}} \cong \{s = (s_i) \in \mathbb{Z}_2^r : \prod_i (s_i)^{n_i} = 1\} / \langle (-1, \dots, -1) \rangle$$

which is a subgroup of $\mathcal{S}_{\underline{\phi}}^{\Sigma_0}$ of index 1 or 2. Similarly, we denote the right hand side by $\mathcal{S}_{\phi}$.

Let us define the characters of $\mathbb{Z}_2^r / \langle (-1, \dots, -1) \rangle$ to be $\mathbb{Z}_2$ -valued functions $\varepsilon = (\varepsilon_i) \in \mathbb{Z}_2^r$ such that $\prod_i \varepsilon_i = 1$. Moreover, for $s \in \mathbb{Z}_2^r / \langle (-1, \dots, -1) \rangle$, we define $\varepsilon(s) = \prod_i (\varepsilon_i * s_i)$, where

$$\varepsilon_i * s_i = \begin{cases} -1, & \text{if } \varepsilon_i = s_i = -1 \\ 1, & \text{otherwise.} \end{cases}$$

So

$$\widehat{\mathcal{S}}_{\phi}^{\Sigma_0} = \{\varepsilon = (\varepsilon_i) \in \mathbb{Z}_2^r : \prod_i \varepsilon_i = 1\}.$$

In particular, when G is special even orthogonal, we define $\varepsilon_0 = (\varepsilon_{0,i}) \in \widehat{\mathcal{S}}_{\phi}^{\Sigma_0}$ satisfying $\varepsilon_{0,i} = 1$ if n_i is even, and $\varepsilon_{0,i} = -1$ if n_i is odd, then

$$\widehat{\mathcal{S}}_{\phi} = \{\varepsilon = (\varepsilon_i) \in \mathbb{Z}_2^r : \prod_i \varepsilon_i = 1\} / \langle \varepsilon_0 \rangle.$$

In general, let $\varepsilon_0 = 1$ if G is not special even orthogonal.

Now we can formulate the theorem for parametrizing supercuspidal representations inside tempered Arthur packets.

Theorem 3.3 (Mœglin [Mœg11], Theorem 1.5.1). *The Σ_0 -orbits of irreducible supercuspidal representations of $G(F)$ can be parametrized by $\phi \in \bar{\Phi}_2(G)$ and $\bar{\varepsilon} \in \widehat{\mathcal{S}}_{\phi}$ satisfying the following properties:*

- (1) if $(\rho, a) \in Jord(\phi)$, then $(\rho, a - 2) \in Jord(\phi)$ as long as $a - 2 > 0$;
- (2) if $(\rho, a), (\rho, a - 2) \in Jord(\phi)$, then $\varepsilon(\rho, a)\varepsilon(\rho, a - 2) = -1$;
- (3) if $(\rho, 2) \in Jord(\phi)$, then $\varepsilon(\rho, 2) = -1$.

The proof that we are going to give makes use of the (twisted) endoscopic character identities and explicit computation of Jacquet modules. So we will first review these two subjects in the next two sections.

4. ENDOSCOPY

The endoscopy theory can be stated for any connected reductive groups, but here we will mainly consider the case when G is a quasisplit symplectic group or special orthogonal group.

Suppose $\phi \in \Phi_2(G)$ and $s \in \bar{S}_{\underline{\phi}} = \mathcal{S}_{\underline{\phi}}$. In our case, there is a quasisplit reductive group H with the property that

$$\widehat{H} \cong \text{Cent}(s, \widehat{G})^0,$$

and the isomorphism extends to an embedding

$$\xi : {}^L H \rightarrow {}^L G$$

such that $\xi({}^L H) \subseteq \text{Cent}(s, {}^L G)$ and $\underline{\phi}$ factors through ${}^L H$. Hence we get a parameter $\phi_H \in \Phi_{bdd}(H)$. In fact it is easy to show $\bar{S}_{\underline{\phi}_H}$ is also finite, so $\phi_H \in \Phi_2(H)$. We say $(H, \underline{\phi}_H)$ corresponds to $(\underline{\phi}, s)$ through ξ , and denote this relation by $(H, \underline{\phi}_H) \rightarrow (\underline{\phi}, s)$. Such H is called an **elliptic endoscopic group** of G .

Example 4.1. (1) If $G = Sp(2n)$, then ${}^L G = SO(2n+1, \mathbb{C})$. For $\phi \in \bar{\Phi}_2(G)$, let us write

$$\phi = \phi_1 \oplus \cdots \oplus \phi_r$$

as in (2.1). Then $\mathcal{S}_\phi = \mathbb{Z}_2^r / \langle -1, \dots, -1 \rangle$, and for any $s = (s_i) \in \mathcal{S}_\phi$, it gives a partition on $Jord(\phi)$, i.e.,

$$\phi = (\oplus_{s_i=1} \phi_i) \oplus (\oplus_{s_j=-1} \phi_j).$$

Without loss of generality, let us assume

$$\sum_{s_i=1} n_i = 2n_I + 1 = N_I \text{ and } \sum_{s_j=-1} n_j = 2n_{II} = N_{II}.$$

Define

$$\eta_I = \eta_{II} = \prod_{s_j=-1} \eta_j,$$

where η_j is a quadratic character given by the central character of π_{ϕ_j} . Let

$$G_I = Sp(2n_I) \text{ and } G_{II} = SO(2n_{II}, \eta_{II}).$$

Then we have

$$H = G_I \times G_{II} \text{ and } {}^L H = (\widehat{G}_I \times \widehat{G}_{II}) \rtimes \Gamma_{E_{II}/F},$$

where E_{II} is a quadratic extension of F associated with η_{II} . Let

$$\xi_i : {}^L G_i \hookrightarrow GL(N_i, \mathbb{C})$$

be the natural embedding for $i = I, II$. Then

$$\xi := (\xi_I \otimes \eta_I) \oplus \xi_{II}$$

factors through ${}^L G$ and defines an embedding ${}^L H \hookrightarrow {}^L G$. Let

$$\phi_I := (\oplus_{s_i=1} \phi_i) \otimes \eta_I \in \bar{\Phi}_2(G_I)$$

and

$$\phi_{II} := \oplus_{s_j=-1} \phi_j \in \bar{\Phi}_2(G_{II}).$$

Then

$$\phi_H = \phi_I \times \phi_{II}.$$

(2) If $G = SO(2n+1)$, then ${}^L G = Sp(2n, \mathbb{C})$. For $\phi \in \bar{\Phi}_2(G)$, let us write

$$\phi = \phi_1 \oplus \cdots \oplus \phi_r$$

as in (2.1). Then $\mathcal{S}_\phi = \mathbb{Z}_2^r / \langle -1, \dots, -1 \rangle$, and for any $s = (s_i) \in \mathcal{S}_\phi$, it gives a partition on $Jord(\phi)$, i.e.,

$$\phi = (\oplus_{s_i=1} \phi_i) \oplus (\oplus_{s_j=-1} \phi_j).$$

We can assume

$$\sum_{s_i=1} n_i = 2n_I = N_I \text{ and } \sum_{s_j=-1} n_j = 2n_{II} = N_{II}.$$

Let

$$G_I = SO(2n_I + 1) \text{ and } G_{II} = SO(2n_{II} + 1).$$

Then we have

$$H = G_I \times G_{II} \text{ and } {}^L H = \widehat{G}_I \times \widehat{G}_{II}$$

Let

$$\xi_i : {}^L G_i \hookrightarrow GL(N_i, \mathbb{C})$$

be the natural embedding for $i = I, II$. Then

$$\xi := \xi_I \oplus \xi_{II}$$

factors through ${}^L G$ and defines an embedding ${}^L H \hookrightarrow {}^L G$. Let

$$\phi_I := (\oplus_{s_i=1} \phi_i) \in \bar{\Phi}_2(G_I)$$

and

$$\phi_{II} := \oplus_{s_j=-1} \phi_j \in \bar{\Phi}_2(G_{II}).$$

Then

$$\phi_H = \phi_I \times \phi_{II}.$$

- (3) If $G = SO(2n, \eta)$, then ${}^L G = SO(2n, \mathbb{C}) \rtimes \Gamma_{E/F}$. For $\phi \in \bar{\Phi}_2(G)$, let us write

$$\phi = \phi_1 \oplus \cdots \oplus \phi_r$$

as in (2.1). Then $\mathcal{S}_\phi^{\Sigma_0} = \mathbb{Z}_2^r / \langle -1, \dots, -1 \rangle$, and for any $s = (s_i) \in \mathcal{S}_\phi \subseteq \mathcal{S}_\phi^{\Sigma_0}$, it gives a partition on $Jord(\phi)$, i.e.,

$$\phi = (\oplus_{s_i=1} \phi_i) \oplus (\oplus_{s_j=-1} \phi_j).$$

By our description of $\mathcal{S}_\phi$, we can assume

$$\sum_{s_i=1} n_i = 2n_I = N_I \text{ and } \sum_{s_j=-1} n_j = 2n_{II} = N_{II}.$$

Define

$$\eta_I = \eta_{II} \eta \text{ and } \eta_{II} = \prod_{s_j=-1} \eta_j,$$

where η_j is a quadratic character given by the central character of π_{ϕ_j} . We also denote by E_i the quadratic extension of F associated with η_i for $i = I, II$. Let

$$G_I = SO(2n_I, \eta_I) \text{ and } G_{II} = SO(2n_{II}, \eta_{II}),$$

Then we have

$$H = G_I \times G_{II} \text{ and } {}^L H = (\widehat{G}_I \times \widehat{G}_{II}) \rtimes \Gamma_{L/F}$$

where $L = E_I E_{II}$. Let

$$\xi_i : {}^L G_i \hookrightarrow GL(N_i, \mathbb{C})$$

be the natural embedding for $i = I, II$. Then

$$\xi := \xi_I \oplus \xi_{II}$$

factors through ${}^L G$ and defines an embedding ${}^L H \hookrightarrow {}^L G$. Let

$$\phi_I := \oplus_{s_i=1} \phi_i \in \bar{\Phi}_2(G_I)$$

and

$$\phi_{II} := \oplus_{s_j=-1} \phi_j \in \bar{\Phi}_2(G_{II})$$

Then

$$\phi_H = \phi_I \times \phi_{II}.$$

In the examples above, we can define $\bar{\Phi}_2(H) = \bar{\Phi}_2(G_I) \times \bar{\Phi}_2(G_{II})$ (resp. $\bar{\Phi}_{bdd}(H) = \bar{\Phi}_{bdd}(G_I) \times \bar{\Phi}_{bdd}(G_{II})$), then $\phi_H \in \bar{\Phi}_2(H)$. For $s \in \mathcal{S}_\phi$, we still say (H, ϕ_H) corresponds to (ϕ, s) , and denote this relation again by $(H, \phi_H) \rightarrow (\phi, s)$.

In part (3), it is possible to also choose $s \in \mathcal{S}_\phi^{\Sigma_0}$ but not in $\mathcal{S}_\phi$, and then we get a partition on $Jord(\phi)$, i.e.,

$$\phi = (\oplus_{s_i=1} \phi_i) \oplus (\oplus_{s_j=-1} \phi_j),$$

so that

$$\sum_{s_i=1} n_i = 2n_I + 1 \text{ and } \sum_{s_j=-1} n_j = 2n_{II} + 1.$$

Define

$$\eta_I = \eta_{II} \eta \text{ and } \eta_{II} = \prod_{s_j=-1} \eta_j,$$

where η_j is a quadratic character given by the central character of π_{ϕ_j} . Let

$$G_I = Sp(2n_I) \text{ and } G_{II} = Sp(2n_{II})$$

Then

$$\phi_I := (\oplus_{s_i=1} \phi_i) \otimes \eta_I \in \bar{\Phi}_2(G_I)$$

and

$$\phi_{II} := (\oplus_{s_j=-1} \phi_j) \otimes \eta_{II} \in \bar{\Phi}_2(G_{II}).$$

We can take

$$H = G_I \times G_{II} \text{ and } {}^L H = \widehat{G}_I \times \widehat{G}_{II}.$$

In this case, H is called a **twisted elliptic endoscopic group** of G . Let

$$\xi_i : {}^L G_i \hookrightarrow GL(N_i, \mathbb{C})$$

be the natural embedding for $i = I, II$. Then

$$\xi := (\xi_I \otimes \eta_I) \oplus (\xi_{II} \otimes \eta_{II})$$

factors through ${}^L G$ and defines an embedding ${}^L H \hookrightarrow {}^L G$. Let

$$\phi_H = \phi_I \times \phi_{II}.$$

We say (H, ϕ_H) corresponds to (ϕ, s) through ξ , and write $(H, \phi_H) \rightarrow (\phi, s)$.

In this paper, we also want to consider the twisted elliptic endoscopic groups of $GL(N)$, but we will only need the simplest case here. Recall for $\phi \in \Phi_{bdd}(G)$, we can view $\underline{\phi}$ as a self-dual N -dimensional representation through the natural embedding

$$\xi_N : {}^L G \rightarrow GL(N, \mathbb{C}),$$

and in this way we get a self-dual parameter for $GL(N)$. We fix an outer automorphism θ_N of $GL(N)$ preserving an F -splitting, and let $\widehat{\theta}_N$ be the dual automorphism on $GL(N, \mathbb{C})$, then

$$\xi_N({}^L G) \subseteq \text{Cent}(s, GL(N, \mathbb{C})) \text{ and } \widehat{G} = \text{Cent}(s, GL(N, \mathbb{C}))^0$$

for some $s \in GL(N, \mathbb{C}) \rtimes \widehat{\theta}_N$. So we call G a twisted elliptic endoscopic group of $GL(N)$.

What lies in the heart of the endoscopy theory is a transfer map on the spaces of smooth compactly supported functions from G to its (twisted) elliptic endoscopic group H (similarly from $GL(N)$ to its twisted elliptic endoscopic group G). The existence of the transfer map is quite deep, and it was conjectured by Langlands, Shelstad and Kottwitz. In a series of papers Waldspurger [Wal95] [Wal97] [Wal06] [Wal08] is able to reduce it to the **Fundamental Lemma** for Lie algebras over function fields. Finally it is in this particular form of the fundamental lemma, Ngo [Ngô10] gave his celebrated proof. Let us denote such transfers by

$$(4.1) \quad C_c^\infty(G(F)) \longrightarrow C_c^\infty(H(F))$$

$$f \longrightarrow f^H$$

and similarly

$$(4.2) \quad C_c^\infty(GL(N, F)) \longrightarrow C_c^\infty(G(F))$$

$$f \longrightarrow f^G$$

We should point out these transfer maps are only well defined after we pass to the space of (twisted) **orbital integrals** on the source and the space of **stable orbital integrals** on the target. Note the space of (twisted) (resp. stable) orbital integrals are dual to the space of (twisted) (resp. stable) invariant distributions on $G(F)$, i.e. one can view the (twisted) (resp. stable) invariant distributions of $G(F)$ as linear functionals of the space of (twisted) (resp. stable) orbital integrals. So dual to these transfer maps, the stable invariant distributions on $H(F)$ (resp. $G(F)$) will map to the (twisted) invariant distributions on $G(F)$ (resp. $GL(N, F)$). We call this map the (twisted) **spectral endoscopic transfer**.

If π is an irreducible smooth representation of $G(F)$, then it defines an invariant distribution on $G(F)$ by the trace of

$$\pi(f) = \int_{G(F)} f(g) \pi(g) dg$$

for $f \in C_c^\infty(G(F))$. We call this the character of π and denote it by $f_G(\pi)$. For any irreducible representation π^{Σ_0} of $G^{\Sigma_0}(F)$, which contains π in its restriction to $G(F)$, we define a twisted invariant distribution on $G(F)$ by the trace of

$$\pi^{\Sigma_0}(f) = \int_{G(F) \rtimes \theta_0} f(g) \pi^{\Sigma_0}(g) dg$$

for $f \in C_c^\infty(G(F) \rtimes \theta_0)$. We call this the twisted character of π , and denote it by $f_G(\pi^{\Sigma_0})$. We can also define the twisted characters for $GL(N)$ similarly, but we will write it in a slightly different way. Let π be a self-dual irreducible smooth representation of $GL(N, F)$, we can define a twisted invariant distribution on $GL(N, F)$ by taking the trace of

$$\pi(f) \circ A_\pi(\theta_N)$$

for $f \in C_c^\infty(GL(N, F))$, where $A_\pi(\theta_N)$ is an intertwining operator between π and π^{θ_N} . We call this the twisted character of π and denote it by $f_{N^\theta}(\pi)$.

Since the (twisted) elliptic endoscopic groups H in our case are all products of quasisplit symplectic and special orthogonal groups, we can define a group of automorphisms of H by taking the product of Σ_0 on each factor, and we denote this group again by Σ_0 . Let $\bar{\mathcal{H}}(G)$ (resp. $\bar{\mathcal{H}}(H)$) be the subspace of Σ_0 -invariant functions in $C_c^\infty(G(F))$ (resp. $C_c^\infty(H(F))$). Then it follows from a simple property of the transfer map (which we will not explain here) that we can restrict both (4.1) and (4.2) to $\bar{\mathcal{H}}(G)$ and $\bar{\mathcal{H}}(H)$. Now we are ready to state a more precise version of Theorem 2.2.

Theorem 4.2 (Arthur). (1) Suppose $\phi \in \bar{\Phi}_2(G)$, the sum of characters in $\bar{\Pi}_\phi$

$$f(\phi) = \sum_{[\pi] \in \bar{\Pi}_\phi} f_G(\pi)$$

defines a stable invariant distribution for $f \in \bar{\mathcal{H}}(G)$. Moreover it is **uniquely** determined by π_ϕ through

$$(4.3) \quad f^G(\phi) = f_{N^\theta}(\pi_\phi), \quad f \in C_c^\infty(GL(N))$$

after we normalize the Haar measures on $G(F)$ and $GL(N, F)$ in a compatible way.

- (2) Suppose $\phi \in \bar{\Phi}_2(G)$, and $(H, \phi_H) \rightarrow (\phi, s)$ for $s \in \mathcal{S}_\phi$. If we define a stable invariant distribution $f(\phi_H)$ for $\bar{\mathcal{H}}(H)$ as in (1), then after we normalize the Haar measures on $G(F)$ and $H(F)$ in a compatible way the following identity holds

$$(4.4) \quad f^H(\phi_H) = \sum_{[\pi] \in \bar{\Pi}_\phi} \langle s, \pi \rangle f_G(\pi) \quad f \in \bar{\mathcal{H}}(G)$$

where

$$\langle \cdot, \pi \rangle := \langle \cdot, \pi \rangle_\phi$$

under the isomorphism $\mathcal{S}_\phi \cong \mathcal{S}_{\underline{\phi}}$.

Remark 4.3. Although we only state the theorem for discrete parameters, these statements are also true for tempered parameters (once we extend the definition $(H, \phi_H) \rightarrow (\phi, s)$ appropriately). The two identities (4.3) and (4.4) are the ones we call (twisted) endoscopic character identities in the end of Section 3, and they are also often referred to as (twisted) character relations. There are some ambiguities that we need to clarify in such identities. On one hand, in the definition of $f_{N^\theta}(\pi_\phi)$ we need to choose a normalization of the intertwining operator $A_{\pi_\phi}(\theta_N)$. In this theorem, we require $A_{\pi_\phi}(\theta_N)$ to fix some Whittaker functional for π_ϕ . On the other hand, in the definition of the transfer maps there is also a normalization issue. To resolve that, we need to fix certain (resp. θ_N -stable) Whittaker datum for G (resp. $GL(N)$), and we will take the so-called Whittaker normalization on the transfer maps. Finally, the stable invariant distribution $f(\phi)$ for $f \in \bar{\mathcal{H}}(G)$ is uniquely determined by π_ϕ for the transfer map (4.2) is surjective onto the space of Σ_0 -invariant stable orbital integrals of $G(F)$.

When G is special even orthogonal, we have an additional character identity. To state it, we need to identify $C_c^\infty(G(F) \rtimes \theta_0)$ with $C_c^\infty(G(F))$ by sending $g \rtimes \theta_0$ to g , so the twisted transfer map on $C_c^\infty(G(F))$ can also be translated to $C_c^\infty(G(F) \rtimes \theta_0)$.

Theorem 4.4 (Arthur). *Suppose $\phi \in \bar{\Phi}_2(G)$, and $(H, \phi_H) \rightarrow (\phi, s)$ for $s \in \mathcal{S}_\phi^{\Sigma_0}$ but not in $\mathcal{S}_\phi$. Then after we normalize the Haar measures on $G(F)$ and $H(F)$ in a compatible way the following identity holds*

$$(4.5) \quad f^H(\phi_H) = \sum_{[\pi] \in \bar{\Pi}_\phi} \langle s, \pi^{\Sigma_0} \rangle f_G(\pi^{\Sigma_0}), \quad f \in C_c^\infty(G(F) \rtimes \theta_0)$$

where $\pi^{\Sigma_0}|_G = \pi$ and

$$\langle \cdot, \pi^{\Sigma_0} \rangle := \langle \cdot, \pi^{\Sigma_0} \rangle_\phi$$

under the isomorphism $\mathcal{S}_\phi^{\Sigma_0} \cong \mathcal{S}_{\underline{\phi}}^{\Sigma_0}$.

Again this theorem also holds for $\phi \in \bar{\Phi}(G)$ (once we extend the definition $(H, \phi_H) \rightarrow (\phi, s)$ appropriately), and we have taken the Whittaker normalization on the transfer maps with respect to the fixed Σ_0 -stable Whittaker datum in Theorem 2.3. We will only need this theorem in Section 9.

5. JACQUET MODULES

First let us assume G is any connected reductive group over F , and let $\text{Rep}(G)$ be the category of finite-length smooth representations of G . If M is the Levi component of a parabolic subgroup P of G , then the normalized parabolic induction defines a functor from $\text{Rep}(M)$ to $\text{Rep}(G)$. The normalized Jacquet module is its left adjoint functor, i.e.,

$$(5.1) \quad \text{Hom}_M(\text{Jac}_P \pi, \sigma) \cong \text{Hom}_G(\pi, \text{Ind}_P^G \sigma),$$

for $\pi \in \text{Rep}(G)$ and $\sigma \in \text{Rep}(M)$. This relation (5.1) is usually referred to as **Frobenius reciprocity**. One can see easily from (5.1) and Theorem 1.1 that $\pi \in \text{Rep}(G)$ is supercuspidal if and only if $\text{Jac}_P \pi = 0$ for all standard parabolic subgroups P of G . In fact this is one of the equivalent definitions of supercuspidal representations. The next two lemmas state some general facts about Jacquet modules, and we refer the interested readers to [MT02, Section 3] for their proofs.

Lemma 5.1. *Suppose $\pi \in \text{Rep}(G)$ is irreducible, and σ is an irreducible supercuspidal constituent of $\text{Jac}_P \pi$, then there is an inclusion*

$$\pi \hookrightarrow \text{Ind}_P^G \sigma.$$

Lemma 5.2. *Suppose $\pi \in \text{Rep}(G)$ is irreducible, and $M = M_1 \times M_2$. Let $\tau_1 \in \text{Rep}(M_1)$ be irreducible and $\tau_2 \in \text{Rep}(M_2)$. If*

$$\pi \hookrightarrow \text{Ind}_P^G(\tau_1 \otimes \tau_2),$$

then there exists an irreducible constituent τ'_2 in τ_2 such that

$$\pi \hookrightarrow \text{Ind}_P^G(\tau_1 \otimes \tau'_2).$$

Now let us restrict to the case when G is a quasisplit symplectic or special orthogonal group. We would like to define a modified Jacquet functor. For this we first fix a unitary irreducible supercuspidal representation ρ of $GL(d_\rho, F)$, and we assume $M = GL(d_\rho) \times G_-$ is the Levi component of a standard maximal parabolic subgroup P of G . In case $G_- = 1$ and G is special even orthogonal, we require P to be contained in the standard parabolic subgroup of $GL(2n)$. Then for $\pi \in \text{Rep}(G)$,

$$s.s.\text{Jac}_P(\pi) = \bigoplus_i \tau_i \otimes \sigma_i,$$

where $\tau_i \in \text{Rep}(GL(d_\rho))$ and $\sigma_i \in \text{Rep}(G_-)$, both of which are irreducible. We define $\text{Jac}_x \pi$ for any real number x to be

$$\text{Jac}_x(\pi) = \bigoplus_{\tau_i = \rho||^x} \sigma_i.$$

Note unlike $\text{Jac}_P \pi$, in our definition $\text{Jac}_x \pi$ is always semisimple. If we have an ordered sequence of real numbers $\{x_1, \dots, x_s\}$, we can define

$$\text{Jac}_{x_1, \dots, x_s} \pi = \text{Jac}_{x_s} \circ \dots \circ \text{Jac}_{x_1} \pi.$$

It is not hard to see Jac_x can be defined for $GL(n)$ in a similar way by replacing G_- by $GL(n_-)$. Furthermore, we can define Jac_x^{op} analogous to Jac_x but with respect to $\rho^\vee$ and the standard Levi subgroup $GL(n_-) \times GL(d_{\rho^\vee})$. So let us define $\text{Jac}_x^\theta = \text{Jac}_x \circ \text{Jac}_{-x}^{op}$ for $GL(n)$. Next we want to give some properties of this modified Jacquet functor.

Lemma 5.3. *If $\pi \in \text{Rep}(G)$ is irreducible, and $\text{Jac}_{x, \dots, y} \pi = \sigma$ for $\sigma \in \text{Rep}(G_-)$. Then there exists an irreducible constituent σ' in σ so that we get an inclusion*

$$\pi \hookrightarrow \rho||^x \times \dots \times \rho||^y \rtimes \sigma'.$$

Proof. By Theorem 1.1, there exists a standard parabolic subgroup P_- of G_- with an irreducible supercuspidal representation π_{M_-} on the Levi component M_- such that there is a nontrivial equivariant homomorphism from σ to $\text{Ind}_{P_-}^{G_-} \pi_{M_-}$. Then by Frobenius reciprocity, π_{M_-} is in $s.s.\text{Jac}_{P_-} \sigma$. In particular, we can take $M = GL(d_\rho) \times \dots \times GL(d_\rho) \times M_-$ with P being the corresponding standard parabolic subgroup of G , and take $\pi_M = \rho||^x \otimes \dots \otimes \rho||^y \otimes \pi_{M_-}$ to be an irreducible supercuspidal representation of M . Then π_M is in $s.s.\text{Jac}_P \pi$. By Lemma 5.1, we know

$$\pi \hookrightarrow \rho||^x \times \dots \times \rho||^y \rtimes \text{Ind}_{P_-}^{G_-}(\pi_{M_-}).$$

So by Lemma 5.2 there exists an irreducible constituent σ' in $\text{Ind}_{M_-}^{G_-}(\pi_{M_-})$ such that

$$\pi \hookrightarrow \rho||^x \times \dots \times \rho||^y \rtimes \sigma'.$$

Finally by Frobenius reciprocity again, we know σ' is in $\text{Jac}_{x, \dots, y} \pi = \sigma$. This finishes the proof. $\square$

As a special case of this lemma, we have the following corollary.

Corollary 5.4. *If $\pi \in \text{Rep}(G)$ is irreducible, and $\text{Jac}_{x, \dots, y} \pi = \sigma$ for $\sigma \in \text{Rep}(G_-)$, which is also irreducible. Then there is an inclusion*

$$\pi \hookrightarrow \rho||^x \times \dots \times \rho||^y \rtimes \sigma.$$

Remark 5.5. Lemma 5.3 and Corollary 5.4 are also valid in the case of general linear groups, and the proofs are the same.

Lemma 5.6. *If $\pi \in \text{Rep}(G)$ and $|x - y| \neq 1$, then $\text{Jac}_{x,y}\pi = \text{Jac}_{y,x}\pi$.*

Proof. We take the standard parabolic subgroup $P = MN$ of G with $M = GL(2d_\rho) \times G_-$. If

$$\text{s.s. Jac}_P \pi = \bigoplus_i \tau_i \otimes \sigma_i,$$

then σ_i is in $\text{Jac}_{x,y}\pi$ if and only if $\text{Jac}_{x,y}\tau_i \neq 0$. Let us assume $\text{Jac}_{x,y}\tau_i \neq 0$, by Corollary 5.4 (also see Remark 5.5) we have $\tau_i \hookrightarrow \rho||^x \times \rho||^y$. Since $|x - y| \neq 1$, $\rho||^x \times \rho||^y \cong \rho||^y \times \rho||^x$ is irreducible (see Appendix B), so we must have $\tau_i \cong \rho||^x \times \rho||^y$. Hence

$$\text{Jac}_{x,y}\pi = \bigoplus_{\tau_i \cong \rho||^x \times \rho||^y} (\text{Jac}_{x,y}\tau_i) \otimes \sigma_i.$$

By the same argument, we have

$$\text{Jac}_{y,x}\pi = \bigoplus_{\tau_i \cong \rho||^y \times \rho||^x} (\text{Jac}_{y,x}\tau_i) \otimes \sigma_i.$$

Therefore, $\text{Jac}_{x,y}\pi = \text{Jac}_{y,x}\pi$. □

Lemma 5.7. *Suppose $\pi \in \text{Rep}(GL(d_\rho(|a - b| + 1)))$ is an irreducible constituent of*

$$\rho||^a \times \cdots \times \rho||^b$$

for a segment $\{a, \dots, b\}$, and $\text{Jac}_x\pi = 0$ unless $x = a$, then $\pi = \langle a, \dots, b \rangle$.

Proof. It is clear that $\text{Jac}_x\pi = 0$ unless $x \in \{a, \dots, b\}$. Suppose $\{a, \dots, y\} \subseteq \{a, \dots, b\}$ is the longest segments such that

$$\text{Jac}_{a, \dots, y}\pi \neq 0.$$

If $y \neq b$, then we can find $z \in \{a, \dots, b\} \setminus \{a\}$ such that $|x - z| > 1$ for all $x \in \{a, \dots, y\}$ and $\text{Jac}_{a, \dots, y, z}\pi \neq 0$. By Lemma 5.6,

$$\text{Jac}_{z, a, \dots, y}\pi = \text{Jac}_{a, \dots, y, z}\pi \neq 0.$$

This means $\text{Jac}_z\pi \neq 0$, and we get a contradiction. So we can only have $y = b$, and by Corollary 5.4 we have

$$\pi \hookrightarrow \rho||^a \times \cdots \times \rho||^b.$$

Hence $\pi = \langle a, \dots, b \rangle$. □

There are some explicit formulas for computing the Jacquet modules in the case of classical groups and general linear groups (see [MT02, Section 1]), and we want to recall some of them here. We will use “ $\stackrel{\text{s.s.}}{=}$ ” for equality after semisimplification.

For $GL(n)$, we know the irreducible discrete series representations are given by

$$\text{St}(\rho', a) = \langle \rho'; \frac{a-1}{2}, \dots, -\frac{a-1}{2} \rangle.$$

More generally we have irreducible representations $\langle \rho'; \zeta a, \dots, \zeta b \rangle$ attached to any decreasing segment $\{a, \dots, b\}$ (cf. Section 1) for $\zeta = \pm 1$. If we fix ρ as before, then we have the following formulas for their Jacquet modules.

$$(5.2) \quad \text{Jac}_x \langle \rho'; \zeta a, \dots, \zeta b \rangle = \begin{cases} \langle \rho'; \zeta(a-1), \dots, \zeta b \rangle, & \text{if } x = \zeta a \text{ and } \rho' \cong \rho, \\ 0, & \text{otherwise;} \end{cases}$$

and

$$(5.3) \quad \text{Jac}_x^{\text{op}} \langle \rho'; \zeta a, \dots, \zeta b \rangle = \begin{cases} \langle \rho'; \zeta a, \dots, \zeta(b+1) \rangle, & \text{if } x = \zeta b \text{ and } \rho' \cong \rho^\vee, \\ 0, & \text{otherwise.} \end{cases}$$

If $\pi_i \in \text{Rep}(GL(n_i))$ for $i = 1$ or 2 , we have

$$\text{Jac}_x(\pi_1 \times \pi_2) \stackrel{s.s.}{=} (\text{Jac}_x \pi_1) \times \pi_2 \oplus \pi_1 \times (\text{Jac}_x \pi_2),$$

and

$$\text{Jac}_x^{op}(\pi_1 \times \pi_2) \stackrel{s.s.}{=} (\text{Jac}_x^{op} \pi_1) \times \pi_2 \oplus \pi_1 \times (\text{Jac}_x^{op} \pi_2).$$

Suppose $\pi \in \text{Rep}(G)$ and $\tau \in \text{Rep}(GL(d))$. If G is symplectic or special odd orthogonal, then

$$\text{Jac}_x(\tau \rtimes \pi) \stackrel{s.s.}{=} (\text{Jac}_x \tau) \rtimes \pi \oplus (\text{Jac}_x^{op} \tau) \rtimes \pi \oplus \tau \rtimes \text{Jac}_x \pi.$$

If $G = SO(2n, \eta)$, the situation is more complicated, and we would like to divide it into three cases.

(1) When $n \neq d_\rho$ or 0 ,

$$\text{Jac}_x(\tau \rtimes \pi) \stackrel{s.s.}{=} \tau \rtimes \text{Jac}_x \pi \oplus \begin{cases} (\text{Jac}_x \tau) \rtimes \pi \oplus (\text{Jac}_x^{op} \tau) \rtimes \pi & \text{if } d_\rho \text{ is even} \\ (\text{Jac}_x \tau) \rtimes \pi \oplus (\text{Jac}_x^{op} \tau) \rtimes \pi^{\theta_0} & \text{if } d_\rho \text{ is odd} \end{cases}$$

(2) When $n = d_\rho$,

$$\text{Jac}_x(\tau \rtimes \pi) \stackrel{s.s.}{=} \tau \rtimes \text{Jac}_x \pi \oplus (\tau \rtimes \text{Jac}_x \pi^{\theta_0})^{\theta_0} \oplus \begin{cases} (\text{Jac}_x \tau) \rtimes \pi \oplus (\text{Jac}_x^{op} \tau) \rtimes \pi & \text{if } d_\rho \text{ is even} \\ (\text{Jac}_x \tau) \rtimes \pi \oplus (\text{Jac}_x^{op} \tau) \rtimes \pi^{\theta_0} & \text{if } d_\rho \text{ is odd} \end{cases}$$

(3) When $n = 0$ and $d \neq d_\rho$,

$$\text{Jac}_x(\tau \rtimes 1) \stackrel{s.s.}{=} \begin{cases} (\text{Jac}_x \tau) \rtimes 1 \oplus (\text{Jac}_x^{op} \tau) \rtimes 1 & \text{if } d_\rho \text{ is even} \\ (\text{Jac}_x \tau) \rtimes 1 \oplus (\text{Jac}_x^{op} \tau \rtimes 1)^{\theta_0} & \text{if } d_\rho \text{ is odd} \end{cases}$$

(4) When $n = 0$ and $d = d_\rho$,

$$\text{Jac}_x(\tau \rtimes 1) \stackrel{s.s.}{=} \begin{cases} (\text{Jac}_x \tau) \rtimes 1 \oplus (\text{Jac}_x^{op} \tau) \rtimes 1 & \text{if } d_\rho \text{ is even} \\ (\text{Jac}_x \tau) \rtimes 1 & \text{if } d_\rho \text{ is odd} \end{cases}$$

The formulas for special even orthogonal groups here are deduced from [Jan06, Theorem 3.4]. At last we define

$$\bar{\text{Jac}}_x = \begin{cases} \text{Jac}_x + \text{Jac}_x \circ \theta_0, & \text{if } G = SO(2n) \text{ and } n = d_\rho \neq 1, \\ \text{Jac}_x, & \text{otherwise.} \end{cases}$$

Let $\bar{\text{Rep}}(G)$ be the category of finite-length representations of $G(F)$ viewed as $\bar{\mathcal{H}}(G)$ -modules. We denote the elements in $\bar{\text{Rep}}(G)$ by $[\pi]$ for $\pi \in \text{Rep}(G)$, and we call $[\pi]$ is irreducible if π is irreducible. For $[\pi] \in \bar{\text{Rep}}(G)$, let us define

$$\tau \rtimes [\pi] := [\tau \rtimes \pi] \text{ and } \bar{\text{Jac}}_x[\pi] := [\bar{\text{Jac}}_x \pi].$$

Then we can combine all cases into the following single formula

$$(5.4) \quad \bar{\text{Jac}}_x(\tau \rtimes [\pi]) \stackrel{s.s.}{=} (\text{Jac}_x \tau) \rtimes [\pi] \oplus (\text{Jac}_x^{op} \tau) \rtimes [\pi] \oplus \tau \rtimes \bar{\text{Jac}}_x[\pi].$$

Finally, we would like to extend the discussion of this section to the category $\text{Rep}(G^{\Sigma_0})$ of finite-length representations of $G^{\Sigma_0}(F)$. Let $P = MN$ be a standard parabolic subgroup of G . If M is θ_0 -stable, we write $M^{\Sigma_0} := M \rtimes \Sigma_0$ and $P^{\Sigma_0} := P \rtimes \Sigma_0$. Otherwise, we let $M^{\Sigma_0} = M$ and $P^{\Sigma_0} = P$. Note when G^{Σ_0} is even orthogonal group, one can define the normalized parabolic induction and normalized Jacquet module in a similar way (see [Ban99, Section 6]). Suppose $\sigma^{\Sigma_0} \in \text{Rep}(M^{\Sigma_0})$, $\pi^{\Sigma_0} \in \text{Rep}(G^{\Sigma_0})$. It follows from the definition that

$$(\text{Jac}_{P^{\Sigma_0}} \pi^{\Sigma_0})|_M = \text{Jac}_P(\pi^{\Sigma_0}|_G).$$

And

$$(\text{Ind}_{P^{\Sigma_0}}^{G^{\Sigma_0}} \sigma^{\Sigma_0})|_G = \text{Ind}_P^G(\sigma^{\Sigma_0}|_M),$$

unless G is special even orthogonal and $M^{\Sigma_0} = M$, in which case

$$(\text{Ind}_{P^{\Sigma_0}}^{G^{\Sigma_0}} \sigma^{\Sigma_0})|_G = \text{Ind}_P^G(\sigma^{\Sigma_0}|_M) \oplus (\text{Ind}_P^G(\sigma^{\Sigma_0}|_M))^{\theta_0}.$$

Let us define

$$\bar{\text{Jac}}_P = \begin{cases} \text{Jac}_P + \text{Jac}_P \circ \theta_0, & \text{if } G = SO(2n) \text{ and } M^{\theta_0} \neq M, \\ \text{Jac}_P, & \text{otherwise.} \end{cases}$$

And

$$\text{Ind}_P^G[\sigma] := [\text{Ind}_P^G \sigma] \text{ and } \bar{\text{Jac}}_P[\pi] := [\bar{\text{Jac}}_P \pi].$$

Then we have

$$(\text{Jac}_{P^{\Sigma_0}} \text{Ind}_{P^{\Sigma_0}}^{G^{\Sigma_0}} \sigma^{\Sigma_0})|_M = \bar{\text{Jac}}_P \text{Ind}_P^G(\sigma^{\Sigma_0}|_M),$$

and

$$[(\text{Ind}_{P^{\Sigma_0}}^{G^{\Sigma_0}} \text{Jac}_{P^{\Sigma_0}} \pi^{\Sigma_0})|_G] = \text{Ind}_P^G \bar{\text{Jac}}_P[\pi^{\Sigma_0}|_G].$$

The Frobenius reciprocity still holds in this case, i.e.,

$$\text{Hom}_{M^{\Sigma_0}}(\text{Jac}_{P^{\Sigma_0}} \pi^{\Sigma_0}, \sigma^{\Sigma_0}) \cong \text{Hom}_{G^{\Sigma_0}}(\pi^{\Sigma_0}, \text{Ind}_{P^{\Sigma_0}}^{G^{\Sigma_0}} \sigma^{\Sigma_0}).$$

Moreover, the results of this section can be stated similarly for representations of $G^{\Sigma_0}(F)$. In particular, for $\tau \in \text{Rep}(GL(d))$, we have

$$(5.5) \quad \text{Jac}_x(\tau \rtimes \pi^{\Sigma_0}) \stackrel{s.s.}{=} (\text{Jac}_x \tau) \rtimes \pi^{\Sigma_0} \oplus (\text{Jac}_{-x}^{\text{op}} \tau) \rtimes \pi^{\Sigma_0} \oplus \tau \rtimes \text{Jac}_x \pi^{\Sigma_0}.$$

6. COMPATIBILITY OF JACQUET MODULES WITH ENDOSCOPIC TRANSFER

As normalized parabolic induction is compatible with endoscopic transfer, the normalized Jacquet module is also compatible with endoscopic transfer. Since the Jacquet module is originally defined on representations, we need to first extend it to the space of finite linear combinations of (twisted) characters, and then to the space of (twisted) invariant distributions (see (C.1)). In particular, this extended Jacquet functor will preserve stability (see (C.3)). If G is any quasisplit connected reductive group over F and θ is an F -automorphism of G preserving an F -splitting, we will denote the space of finite linear combinations of twisted characters of $G(F)$ by $R(G^\theta)$ and denote the space of stable finite linear combinations of characters on $G(F)$ by $R(G)^{st}$. Moreover, let $\widehat{I}(G^\theta)$ be the space of twisted invariant distributions on $G(F)$ and $\widehat{SI}(G)$ be the space of stable invariant distributions on $G(F)$. When $G = GL(N)$, we will simply write $R(N^\theta)$ and $\widehat{I}(N^\theta)$ for the corresponding spaces. In the following discussion we will assume G is a quasisplit symplectic or special orthogonal group.

Suppose H is an elliptic endoscopic group of G , we know from Section 4 that $H = G_I \times G_{II}$, and there is an embedding

$$\xi : {}^L H \hookrightarrow {}^L G,$$

where ${}^L H = \widehat{H} \rtimes \Gamma_{L/F}$ for $L = F, E_{II}$ or $E_I E_{II}$ accordingly. We fix Γ -splittings $(\mathcal{B}_H, \mathcal{T}_H, \{\mathcal{X}_{\alpha_H}\})$ and $(\mathcal{B}_G, \mathcal{T}_G, \{\mathcal{X}_{\alpha_G}\})$ for $\widehat{H}$ and $\widehat{G}$ respectively. By taking certain $\widehat{G}$ -conjugate of ξ , we can assume $\xi(\mathcal{T}_H) = \mathcal{T}_G$ and $\xi(\mathcal{B}_H) \subseteq \mathcal{B}_G$. So we can view the Weyl group $W_H = W(\widehat{H}, \mathcal{T}_H)$ as a subgroup of $W_G = W(\widehat{G}, \mathcal{T}_G)$. We also view ${}^L H$ as a subgroup of ${}^L G$ through ξ .

We fix a standard parabolic subgroup $P = MN$ of G with standard embedding ${}^L P \hookrightarrow {}^L G$. Then there exists a torus $S \subseteq \mathcal{T}_G$ such that ${}^L M = \text{Cent}(S, {}^L G)$. Let $W_M = W(\widehat{M}, \mathcal{T}_G)$. We define

$$W_G(H, M) := \{w \in W_G \mid \text{Cent}(w(S), {}^L H) \rightarrow \Gamma_{L/F} \text{ surjective} \}.$$

For any $w \in W_G(H, M)$, let us take $g \in \widehat{G}$ such that $\text{Int}(g)$ induces w . Since $\text{Cent}(w(S), {}^L H) \rightarrow \Gamma_{L/F}$ is surjective, $g^L P g^{-1} \cap {}^L H$ defines a parabolic subgroup of ${}^L H$ with Levi component $g^L M g^{-1} \cap {}^L H$. So we can choose a standard parabolic subgroup $P'_w = M'_w N'_w$ of H with standard embedding ${}^L P'_w \hookrightarrow {}^L H$ such that ${}^L P'_w$ (resp. ${}^L M'_w$) is $\widehat{H}$ -conjugate to $g^L P g^{-1} \cap {}^L H$ (resp. $g^L M g^{-1} \cap {}^L H$). In fact, M'_w is an endoscopic

group of M (not necessarily elliptic, see [KS99]), and we have an embedding $\xi_{M'_w} : {}^L M'_w \rightarrow {}^L M$ given by the following diagram:

$$\begin{array}{ccccccc} {}^L P'_w & \longleftarrow & {}^L M'_w & \xrightarrow{\xi_{M'_w}} & {}^L M & \longrightarrow & {}^L P \\ \downarrow & & & & & & \downarrow \\ {}^L H & \xrightarrow{\text{Int}(h)} & {}^L H & \xrightarrow{\xi} & {}^L G & \xleftarrow{\text{Int}(g)} & {}^L G \end{array}$$

where $h \in \widehat{H}$ induces an element in W_H . Note the choice of h is unique up to $\widehat{M}'_w$ -conjugation, and so is $\xi_{M'_w}$. If we change g to $h'gm$, where $h' \in \widehat{H}$ induces an element in W_H and $m \in \widehat{M}$ induces an element in W_M , then we still get P'_w , but $\xi_{M'_w}$ changes to $\text{Int}(m^{-1}) \circ \xi_{M'_w}$ up to $\widehat{M}'_w$ -conjugation. Therefore, for any element w in

$$W_H \backslash W_G(H, M) / W_M,$$

we can associate a standard parabolic subgroup $P'_w = M'_w N'_w$ of H and a $\widehat{M}$ -conjugacy class of embedding $\xi_{M'_w} : {}^L M'_w \rightarrow {}^L M$. Moreover, we have the following commutative diagram:

$$(6.1) \quad \begin{array}{ccc} \widehat{SI}(H) & \longrightarrow & \widehat{I}(G) \\ \oplus_w \text{Jac}_{P'_w} \downarrow & & \downarrow \text{Jac}_P \\ \oplus_w \widehat{SI}(M'_w) & \longrightarrow & \widehat{I}(M), \end{array}$$

where the sum is over $W_H \backslash W_G(H, M) / W_M$, and the horizontal maps correspond to the spectral endoscopic transfers with respect to ξ on the top and $\xi_{M'_w}$ on the bottom.

Suppose $M = GL(m) \times G_-$, then the Levi subgroups M'_w of H appearing in (6.1) are of the form $M_I \times M_{II}$, where $M_I \cong GL(m_I) \times G_{I-}$ is a Levi subgroup of G_I and $M_{II} \cong GL(m_{II}) \times G_{II-}$ is a Levi subgroup of G_{II} with $m = m_I + m_{II}$. The spectral endoscopic transfer maps $\widehat{SI}(G_{I-} \times G_{II-})$ to $\widehat{I}(G_-)$, and it also maps $\widehat{SI}(GL(m_I) \times GL(m_{II}))$ to $\widehat{I}(GL(m))$, which is given by parabolic induction. Now we fix a unitary irreducible supercuspidal representation ρ of $GL(d_\rho, F)$, and let $m = d_\rho$. We would like to restrict (6.1) to invariant distributions of $M(F)$ such that on $GL(d_\rho, F)$ they are given by the character of $\rho||^x$, then the relevant Levi subgroups of H will satisfy $m_I = 0$ or $m_{II} = 0$. Since there is no canonical projection from $\widehat{I}(M)$ to such distributions, we will have to restrict (6.1) to spaces of finite linear combinations of characters first.

Let us write

$$M'_{w_I} = GL(d_\rho) \times H_{I-} := GL(d_\rho) \times G_{I-} \times G_{II}$$

and

$$M'_{w_{II}} = GL(d_\rho) \times H_{II-} := GL(d_\rho) \times G_I \times G_{II-}.$$

We also keep the notations in Example 4.1, in particular when G is symplectic, G_I is symplectic and G_{II} is special even orthogonal. Let $\theta_i = \theta_0$ with respect to G_i for $i = I, II$. Then we have the following cases.

- (1) If G is symplectic, then M'_{w_I} , $M'_{w_{II}}$ and $(M'_{w_{II}})^{\theta_{II}}$ are the relevant standard Levi subgroups of H . Note $M'_{w_{II}} = (M'_{w_{II}})^{\theta_{II}}$ if and only if $G_{II-} \neq 1$. We get a modified diagram of (6.1) as follows.

$$(6.2) \quad \begin{array}{ccc} R(H)^{st} & \longrightarrow & R(G) \\ \text{Jac}_x^I \oplus \bar{\text{Jac}}_x \downarrow & & \downarrow \text{Jac}_x \\ R(H_{I-})^{st} \oplus R(H_{II-})^{st} & \longrightarrow & R(G_-), \end{array}$$

where Jac_x^I is with respect to $\rho \otimes \eta_I$.

- (2) If G is special odd orthogonal, then M'_{w_I} , $M'_{w_{II}}$ are the only relevant standard Levi subgroups of H , and we get a modified diagram of (6.1) as follows.

$$(6.3) \quad \begin{array}{ccc} R(H)^{st} & \longrightarrow & R(G) \\ \text{Jac}_x \oplus \text{Jac}_x \downarrow & & \downarrow \text{Jac}_x \\ R(H_{I-})^{st} \oplus R(H_{II-})^{st} & \longrightarrow & R(G_-) \end{array}$$

- (3) If G is special even orthogonal, then M'_{w_I} , $(M'_{w_I})^{\theta_I}$, $M'_{w_{II}}$ and $(M'_{w_{II}})^{\theta_{II}}$ are the relevant standard Levi subgroups of H . Note $M'_{w_i} = (M'_{w_i})^{\theta_i}$ if and only if $G_{i-} \neq 1$ for $i = I, II$. We get a modified diagram of (6.1) as follows.

$$(6.4) \quad \begin{array}{ccc} R(H)^{st} & \longrightarrow & R(G) \\ \bar{\text{Jac}}_x \oplus \bar{\text{Jac}}_x \downarrow & & \downarrow \text{Jac}_x \\ R(H_{I-})^{st} \oplus R(H_{II-})^{st} & \longrightarrow & R(G_-) \end{array}$$

Next we view G as a twisted elliptic endoscopic group of $GL(N)$, and there is an embedding

$$\xi_N : {}^L G \hookrightarrow GL(N, \mathbb{C})$$

where ${}^L G = \widehat{G} \rtimes \Gamma_{L/F}$ for $L = F$ or E accordingly. We also fix a $\widehat{\theta}_N$ -stable Γ -splitting $(\mathcal{B}_N, \mathcal{T}_N, \{\mathcal{X}_{\alpha_N}\})$ of $GL(N, \mathbb{C})$. By taking certain $GL(N, \mathbb{C})$ -conjugate of ξ_N , we can assume $\xi_N(\mathcal{T}_G) = (\mathcal{T}_N^{\widehat{\theta}_N})^0$, $\xi_N(\mathcal{B}_G) \subseteq \mathcal{B}_N$. So we can view the Weyl group $W_G = W(\widehat{G}, \mathcal{T}_G)$ as a subgroup of $W_{N^\theta} := W(GL(N, \mathbb{C}), \mathcal{T}_N)^{\widehat{\theta}_N}$. We also view ${}^L G$ as a subgroup of $GL(N, \mathbb{C})$ through ξ_N .

We fix a standard θ_N -stable parabolic subgroup $P = MN$ of $GL(N)$ with standard embedding ${}^L P \hookrightarrow GL(N, \mathbb{C})$. Then there exists a torus $S \subseteq (\mathcal{T}_N^{\widehat{\theta}_N})^0$ such that ${}^L M = \text{Cent}(S, GL(N, \mathbb{C}))$. Let $W_{M^\theta} = W(\widehat{M}, \mathcal{T}_N)^{\widehat{\theta}_N}$. We define

$$W_{N^\theta}(G, M) := \{w \in W_{N^\theta} \mid \text{Cent}(w(S), {}^L G) \rightarrow \Gamma_{L/F} \text{ surjective} \}.$$

For any $w \in W_{N^\theta}(G, M)$, let us take $g_N \in GL(N, \mathbb{C})$ such that $\text{Int}(g_N)$ induces w . Since $\text{Cent}(w(S), {}^L G) \rightarrow \Gamma_{L/F}$ is surjective, $g_N {}^L P g_N^{-1} \cap {}^L G$ defines a parabolic subgroup of ${}^L G$ with Levi component $g_N {}^L M g_N^{-1} \cap {}^L G$. So we can choose a standard parabolic subgroup $P'_w = M'_w N'_w$ of H with standard embedding ${}^L P'_w \hookrightarrow {}^L G$ such that ${}^L P'_w$ (resp. ${}^L M'_w$) is $\widehat{G}$ -conjugate to $g_N {}^L P g_N^{-1} \cap {}^L G$ (resp. $g_N {}^L M g_N^{-1} \cap {}^L G$). As before, M'_w can be viewed as a twisted endoscopic groups of M (not necessarily elliptic, see [KS99]), and we have an embedding $\xi_{M'_w} : {}^L M'_w \rightarrow {}^L M$ given by the following diagram:

$$\begin{array}{ccccccc} {}^L P'_w & \longleftarrow & {}^L M'_w & \xrightarrow{\xi_{M'_w}} & {}^L M & \longrightarrow & {}^L P \\ \downarrow & & & & & & \downarrow \\ {}^L G & \xrightarrow{\text{Int}(g)} & {}^L G & \xrightarrow{\xi_N} & GL(N, \mathbb{C}) & \xleftarrow{\text{Int}(g_N)} & GL(N, \mathbb{C}) \end{array}$$

where $g \in \widehat{G}$ induces an element in W_G . Note the choice of g is unique up to $\widehat{M}'_w$ -conjugation, and so is $\xi_{M'_w}$. If we change g_N to $g' g_N m$, where $g' \in \widehat{G}$ induces an element in W_G and $m \in \widehat{M}$ induces an element in W_{M^θ} , then we still get P'_w , but $\xi_{M'_w}$ changes to $\text{Int}(m^{-1}) \circ \xi_{M'_w}$ up to $\widehat{M}'_w$ -conjugation. Therefore, for any element w in

$$W_G \backslash W_{N^\theta}(G, M) / W_{M^\theta}$$

we can associate a standard parabolic subgroup $P'_w = M'_w N'_w$ of G and a $\widehat{M}$ -conjugacy class of embedding $\xi_{M'_w} : {}^L M'_w \rightarrow {}^L M$. Moreover, we have the following commutative diagram:

$$(6.5) \quad \begin{array}{ccc} \widehat{SI}(G) & \longrightarrow & \widehat{I}(N^\theta) \\ \oplus_w \text{Jac}_{P'_w} \downarrow & & \downarrow \text{Jac}_P \\ \bigoplus_w \widehat{SI}(M'_w) & \longrightarrow & \widehat{I}(M^\theta). \end{array}$$

where the sum is over $W_G \backslash W_{N^\theta}(G, M) / W_{M^\theta}$, and the horizontal maps correspond to the twisted spectral endoscopic transfers with respect to ξ on the top and $\xi_{M'_w}$ on the bottom.

We again fix a unitary irreducible supercuspidal representation ρ of $GL(d_\rho, F)$, and suppose

$$M = GL(d_\rho) \times GL(N_-) \times GL(d_{\rho^\vee}).$$

Then the standard Levi subgroups of G appearing in (6.5) are

$$M'_w = GL(d_\rho) \times G_-$$

and $(M'_w)^{\theta_0}$. Note $M'_w = (M'_w)^{\theta_0}$ unless G is special even orthogonal and $N_- = 0$. For the purpose of restricting (6.5) to the twisted invariant distributions of $M(F)$ such that on $GL(d_\rho, F) \times GL(d_{\rho^\vee}, F)$ they are given by the twisted character of $\rho ||^x \otimes \rho^\vee ||^{-x}$, we will have to first restrict the diagram to spaces of finite linear combinations of (twisted) characters. Then we can get a modified diagram as follows

$$(6.6) \quad \begin{array}{ccc} R(G)^{st} & \longrightarrow & R(N^\theta) \\ \text{Jac}_x \downarrow & & \downarrow \text{Jac}_x^\theta \\ R(G_-)^{st} & \longrightarrow & R(N_-^\theta). \end{array}$$

At last, when G is special even orthogonal, there is a twisted version of the diagram (6.4), which can be derived as in the case of $GL(N)$ (also see Appendix C for the general case). Here we will only state the result using the notations from Section 4 and (6.4). We assume $G_- \neq 1$.

$$(6.7) \quad \begin{array}{ccc} R(H)^{st} & \longrightarrow & R(G^{\theta_0}) \\ \text{Jac}_x^I \oplus \text{Jac}_x^{II} \downarrow & & \downarrow \text{Jac}_x \\ R(H_{I-})^{st} \oplus R(H_{II-})^{st} & \longrightarrow & R(G_-^{\theta_0}), \end{array}$$

where Jac_x^i is with respect to $\rho \otimes \eta_i$ for $i = I, II$.

Both diagrams (6.1) and (6.5) can be established by using Casselman's formula [Cas77] and its twisted version for relating the (twisted) characters of representations with that of their **unnormalized** Jacquet modules (see [MW06] and [Hir04]). For the convenience of the reader, we will give the proof of the general case in Appendix C. In the next section, we are going to prove Theorem 3.3 by applying (6.6) (resp. (6.2), (6.3) and (6.4)) to the (twisted) endoscopic character identity (4.3) (resp. (4.4)). We will only need (6.7) in Section 9.

7. PROOF OF THEOREM 3.3

In the following sections we will always assume G is a quasisplit symplectic group or special orthogonal group. Before we start the proof, we would like to make explicit the effects of Jacquet modules on the (twisted) endoscopic character identities (4.3) and (4.4). So let us fix a self-dual irreducible unitary supercuspidal representation ρ of $GL(d_\rho, F)$ and a real number x . Let $\phi \in \bar{\Phi}_2(G)$ and we define $\phi_- \in \bar{\Phi}_{bdd}(G_-)$ by its $Jord(\phi_-)$ as follows.

$$Jord(\phi_-) = Jord(\phi) \cup \{(\rho, 2x - 1)\} \setminus \{(\rho, 2x + 1)\},$$

if $(\rho, 2x + 1) \in Jord(\phi)$ and $x > 0$, or $\emptyset$ otherwise. We also set $\pi_{\phi_-} = 0$ if $Jord(\phi_-) = \emptyset$. Note ϕ_- depends on both ρ and x . The following lemma is clear by our explicit formulas (5.2) and (5.3).

Lemma 7.1. $\pi_{\phi_-} = \text{Jac}_x^\theta \pi_\phi$.

So after applying (6.6) to the twisted endoscopic identity (4.3), we have

$$(7.1) \quad f^{G^-} \left(\sum_{[\pi] \in \bar{\Pi}_\phi} \bar{\text{Jac}}_x \pi \right) = f_{N_-^\theta}(\pi_{\phi_-})$$

for $f \in C_c^\infty(GL(N_-), F)$. Since Theorem 4.2 is also valid for all tempered parameters (see Remark 4.3), the left hand side of (7.1) has to be $f^{G^-}(\phi_-)$. Then we have the following result.

Lemma 7.2.

$$(7.2) \quad \bar{\Pi}_{\phi_-} = \bar{\text{Jac}}_x \bar{\Pi}_\phi.$$

In particular,

$$(7.3) \quad \bar{\text{Jac}}_x \bar{\Pi}_\phi = 0 \text{ if } (\rho, 2x+1) \notin \text{Jord}(\phi).$$

Proof. Since the transfer map (4.2) is surjective onto the space of Σ_0 -invariant stable orbital integrals of $G(F)$, we have

$$f(\phi_-) = f \left(\sum_{[\pi] \in \bar{\Pi}_\phi} \bar{\text{Jac}}_x \pi \right) = \sum_{[\pi] \in \bar{\Pi}_\phi} f_G(\bar{\text{Jac}}_x \pi)$$

for $f \in \bar{\mathcal{H}}(G)$. Then the lemma follows from the linear independence of characters of irreducible smooth representations of $G(F)$. $\square$

Suppose $\text{Jord}(\phi_-) \neq \emptyset$, then $x > 0$ and there are two possibilities, i.e., $\phi_- \in \bar{\Phi}_2(G_-)$, or

$$\phi_- = 2\phi_1 \oplus \phi_2 \oplus \cdots \oplus \phi_r \in \bar{\Phi}_{bdd}(G_-)$$

as in (2.1), where $\phi_1 = \rho \otimes [2x-1]$. In the first case we have $(\rho, 2x-1) \notin \text{Jord}(\phi)$. If $x \neq 1/2$, then there is a canonical isomorphism $\mathcal{S}_\phi \cong \mathcal{S}_{\phi_-}$ after identifying $\text{Jord}(\phi)$ with $\text{Jord}(\phi_-)$ by sending $(\rho, 2x+1)$ to $(\rho, 2x-1)$. If $x = 1/2$, we have a projection from $\mathcal{S}_\phi$ to $\mathcal{S}_{\phi_-}$ by restricting $\mathbb{Z}_2$ -valued functions on $\text{Jord}(\phi)$ to $\text{Jord}(\phi_-)$. Hence we get an exact sequence

$$(7.4) \quad 1 \longrightarrow \langle s \rangle \longrightarrow \mathcal{S}_\phi \longrightarrow \mathcal{S}_{\phi_-} \longrightarrow 1,$$

where $s(\cdot) = 1$ over $\text{Jord}(\phi)$ except for $s(\rho, 1/2) = -1$.

In the second case, let $\underline{\phi}_- : L_F \rightarrow {}^L G_-$ be a representative of ϕ_- . We can also identify $\mathcal{S}_{\underline{\phi}_-}$ and its characters $\widehat{\mathcal{S}}_{\underline{\phi}_-}$ with certain quotient spaces of $\mathbb{Z}_2$ -valued functions on $\text{Jord}(\phi_-)$ (forgetting multiplicities), as in the case of discrete parameters. Note

$$\text{Cent}(\underline{\phi}_-, GL(N_-, \mathbb{C})) \cong \underbrace{GL(2, \mathbb{C}) \times \mathbb{C}^\times \times \cdots \times \mathbb{C}^\times}_r,$$

and then

$$\text{Cent}(\underline{\phi}_-, \widehat{G}_-) \cong \{s = (s_i) \in O(2, \mathbb{C}) \times \mathbb{Z}_2^{r-1} : \det(s_1)^{n_1} \cdot \prod_{i \neq 1} (s_i)^{n_i} = 1\}.$$

We write z for the nontrivial central element of $O(2, \mathbb{C})$. Then $\bar{\mathcal{S}}_{\underline{\phi}_-}^{\Sigma_0} \cong O(2, \mathbb{C}) \times \mathbb{Z}_2^{r-1} / \langle z, -1, \dots, -1 \rangle$, and hence $\mathcal{S}_{\underline{\phi}_-}^{\Sigma_0} \cong \mathbb{Z}_2^r / \langle 1, -1, \dots, -1 \rangle$. If G is special even orthogonal,

$$\mathcal{S}_{\underline{\phi}_-} \cong \{s = (s_i) \in \mathbb{Z}_2^r : \prod_i (s_i)^{n_i} = 1\} / \langle 1, -1, \dots, -1 \rangle$$

which is a subgroup of $\mathcal{S}_{\underline{\phi}_-}^{\Sigma_0}$ of index 1 or 2. Let us denote by $\mathcal{S}_{\phi_-}$ (resp. $\mathcal{S}_{\phi_-}^{\Sigma_0}$) the corresponding quotient space of $\mathbb{Z}_2$ -valued functions on $\text{Jord}(\phi_-)$ (forgetting multiplicities) such that $\mathcal{S}_{\phi_-} \cong \mathcal{S}_{\underline{\phi}_-}$ (resp. $\mathcal{S}_{\phi_-}^{\Sigma_0} \cong \mathcal{S}_{\underline{\phi}_-}^{\Sigma_0}$) under these isomorphisms. There is a projection from $\mathcal{S}_\phi \rightarrow \mathcal{S}_{\phi_-}$ (resp. $\mathcal{S}_\phi^{\Sigma_0} \rightarrow \mathcal{S}_{\phi_-}^{\Sigma_0}$) by sending s to

s_- such that $s_-(\cdot) = s(\cdot)$ over $Jord(\phi) \setminus \{(\rho, 2x+1), (\rho, 2x-1)\}$ and $s_-(\rho, 2x-1) = s(\rho, 2x+1)s(\rho, 2x-1)$. Hence there is a short exact sequence

$$(7.5) \quad 1 \longrightarrow \langle s \rangle \longrightarrow \mathcal{S}_\phi \longrightarrow \mathcal{S}_{\phi_-} \longrightarrow 1$$

$$(\text{ resp. } 1 \longrightarrow \langle s \rangle \longrightarrow \mathcal{S}_\phi^{\Sigma_0} \longrightarrow \mathcal{S}_{\phi_-}^{\Sigma_0} \longrightarrow 1)$$

where $s(\cdot) = 1$ over $Jord(\phi)$ except for $s(\rho, 2x+1) = s(\rho, 2x-1) = -1$. For the characters of $\mathcal{S}_{\phi_-}$, we have

$$\widehat{\mathcal{S}_{\phi_-}^{\Sigma_0}} = \{\varepsilon = (\varepsilon_i \in \mathbb{Z}_2^r) : \prod_{i \neq 1} \varepsilon_i = 1\},$$

and if G is special even orthogonal,

$$\widehat{\mathcal{S}_{\phi_-}} = \{\varepsilon = (\varepsilon_i \in \mathbb{Z}_2^r) : \prod_{i \neq 1} \varepsilon_i = 1\} / \langle \varepsilon_0 \rangle,$$

where $\varepsilon_0 = (\varepsilon_{0,i}) \in \widehat{\mathcal{S}_{\phi_-}^{\Sigma_0}}$ satisfies $\varepsilon_{0,i} = 1$ if n_i is even, and $\varepsilon_{0,i} = -1$ if n_i is odd. So ε_0 is trivial when restricted to $\mathcal{S}_{\phi_-}$. In general, let $\varepsilon_0 = 1$ if G is not special even orthogonal.

At last we want to point out in this case ϕ_- factors through $\phi_{M_-} \in \bar{\Phi}_2(M_-)$, for a Levi subgroup $M_- \cong GL(n_1) \times G'$ and $\phi_{M_-} = \phi_1 \times \phi'$ such that

$$\phi' = \phi_2 \oplus \cdots \oplus \phi_r.$$

Since

$$\mathcal{S}_{\phi_{M_-}}^{\Sigma_0} \hookrightarrow \mathcal{S}_{\phi_-}^{\Sigma_0} \text{ and } \mathcal{S}_{\phi_{M_-}}^{\Sigma_0} \cong \mathcal{S}_{\phi'}^{\Sigma_0},$$

we can get an inclusion $\mathcal{S}_{\phi'}^{\Sigma_0} \hookrightarrow \mathcal{S}_{\phi_-}^{\Sigma_0}$, which in fact just extends $s'(\cdot) \in \mathbb{Z}_2^{Jord(\phi')}$ trivially to $Jord(\phi_-)$ (forgetting multiplicities). So on the dual side, there is a projection

$$\widehat{\mathcal{S}_{\phi_-}^{\Sigma_0}} \rightarrow \widehat{\mathcal{S}_{\phi'}^{\Sigma_0}}$$

given by restricting $\varepsilon(\cdot)$ to $Jord(\phi')$. Taking quotient by $\langle \varepsilon_0 \rangle$, we get $\widehat{\mathcal{S}_{\phi_-}} \rightarrow \widehat{\mathcal{S}_{\phi'}}$. It follows from Arthur's theory (i.e., Theorem 2.2, 2.3, 4.2 and 4.4) that

$$(7.6) \quad \bar{\Pi}_{\phi_-} = \pi_{\phi_1} \rtimes \bar{\Pi}_{\phi'} \quad (\text{ resp. } \Pi_{\phi_-}^{\Sigma_0} = \pi_{\phi_1} \rtimes \Pi_{\phi'}^{\Sigma_0})$$

Moreover,

$$(7.7) \quad \pi_{\phi_1} \rtimes \pi(\phi', \varepsilon') = \bigoplus_{\varepsilon' \leftarrow \bar{\varepsilon} \in \widehat{\mathcal{S}_{\phi_-}}} \pi(\phi_-, \bar{\varepsilon})$$

$$\left(\text{ resp. } \pi_{\phi_1} \rtimes \pi^{\Sigma_0}(\phi', \varepsilon') = \bigoplus_{\varepsilon' \leftarrow \varepsilon \in \widehat{\mathcal{S}_{\phi_-}^{\Sigma_0}}} \pi^{\Sigma_0}(\phi_-, \varepsilon) \right).$$

We will need this description of $\bar{\Pi}_{\phi_-}$ (resp. $\Pi_{\phi_-}^{\Sigma_0}$) in Section 8 (resp. Section 9).

In all the above cases, we can canonically identify $\widehat{\mathcal{S}_{\phi_-}}$ (resp. $\widehat{\mathcal{S}_{\phi_-}^{\Sigma_0}}$) with a subgroup of $\widehat{\mathcal{S}_\phi}$ (resp. $\widehat{\mathcal{S}_\phi^{\Sigma_0}}$) of index 1 or 2, so later on we will always view $\varepsilon \in \widehat{\mathcal{S}_{\phi_-}^{\Sigma_0}}$ as functions on $Jord(\phi)$.

Lemma 7.3. *Suppose $\phi \in \bar{\Phi}_2(G)$, and $(\rho, 2x+1) \in Jord(\phi)$.*

- (1) *If $x > 1/2$ and $(\rho, 2x-1) \notin Jord(\phi)$, then $\pi(\phi_-, \bar{\varepsilon}) = \bar{J}ac_x \pi(\phi, \bar{\varepsilon})$ for all $\bar{\varepsilon} \in \widehat{\mathcal{S}_{\phi_-}} \cong \widehat{\mathcal{S}_\phi}$.*
- (2) *If $x > 1/2$ and $(\rho, 2x-1) \in Jord(\phi)$, then $\bar{J}ac_x \pi(\phi, \bar{\varepsilon}) = 0$ unless $\bar{\varepsilon} \in \widehat{\mathcal{S}_{\phi_-}}$, i.e.,*

$$\varepsilon(\rho, 2x+1)\varepsilon(\rho, 2x-1) = 1,$$

in which case $\pi(\phi_-, \bar{\varepsilon}) = \bar{J}ac_x \pi(\phi, \bar{\varepsilon})$.

(3) If $x = 1/2$, then $\bar{J}ac_{1/2}\pi(\phi, \bar{\varepsilon}) = 0$ unless $\bar{\varepsilon} \in \widehat{\mathcal{S}}_{\phi_-}$, i.e.,

$$\varepsilon(\rho, 2) = 1,$$

in which case $\pi(\phi_-, \bar{\varepsilon}) = \bar{J}ac_x\pi(\phi, \bar{\varepsilon})$.

Proof. First we know from (7.2) that $\bar{\Pi}_{\phi_-} = \bar{J}ac_x\bar{\Pi}_{\phi}$, so in particular $\bar{J}ac_x\pi$ do not have common irreducible constituents with each other for $[\pi] \in \bar{\Pi}_{\phi}$. Next for $s \in \mathcal{S}_{\phi}$, suppose $(H, \phi_H) \rightarrow (\phi, s)$, then we have

$$(7.8) \quad f^H(\phi_H) = \sum_{[\pi] \in \bar{\Pi}_{\phi}} \langle s, \pi \rangle f_G(\pi) = \sum_{\varepsilon \in \widehat{\mathcal{S}}_{\phi}} \varepsilon(s) f_G(\pi(\phi, \varepsilon)),$$

for $f \in \bar{\mathcal{H}}(G)$. In the notation of Section 4, we can write

$$H = G_I \times G_{II} \text{ and } \phi_H = \phi_I \times \phi_{II}$$

Let us first assume $(\rho, 2x+1) \notin \text{Jord}(\phi_{II})$. Then $(\rho', 2x+1) \in \text{Jord}(\phi_I)$ for $\rho' = \rho \otimes \eta_I$ if G is symplectic, and $\rho' = \rho$ otherwise. By (7.3),

$$\bar{J}ac_x\bar{\Pi}_{\phi_{II}} = 0.$$

So we let $H_- = H_{I-}$ (see Section 6), and define $\phi_{H_-} = \phi_{I-} \times \phi_{II}$, where

$$\text{Jord}(\phi_{I-}) = \text{Jord}(\phi_I) \cup \{(\rho', 2x-1)\} \setminus \{(\rho', 2x+1)\}.$$

After applying (6.2), (6.3) and (6.4) accordingly to (7.8), we get

$$(7.9) \quad f^{H_-}(\phi_{H_-}) = \sum_{[\pi] \in \bar{\Pi}_{\phi}} \langle s, \pi \rangle f_{G_-}(\bar{J}ac_x\pi) = \sum_{\bar{\varepsilon} \in \widehat{\mathcal{S}}_{\phi}} \bar{\varepsilon}(s) f_{G_-}(\bar{J}ac_x\pi(\phi, \bar{\varepsilon})),$$

for $f \in \bar{\mathcal{H}}(G_-)$. On the other hand note $(H_-, \phi_{H_-}) \rightarrow (\phi_-, s_-)$, where s_- is the image of s under the projection $\mathcal{S}_{\phi} \rightarrow \mathcal{S}_{\phi_-}$, so we have

$$f^{H_-}(\phi_{H_-}) = \sum_{[\pi_-] \in \bar{\Pi}_{\phi_-}} \langle s_-, \pi_- \rangle f_{G_-}(\pi_-) = \sum_{\bar{\varepsilon}' \in \widehat{\mathcal{S}}_{\phi_-}} \bar{\varepsilon}'(s_-) f_{G_-}(\pi(\phi_-, \bar{\varepsilon}')),$$

for $f \in \bar{\mathcal{H}}(G_-)$. Combining this identity with (7.9), we get

$$(7.10) \quad \sum_{\bar{\varepsilon} \in \widehat{\mathcal{S}}_{\phi}} \bar{\varepsilon}(s) f_{G_-}(\bar{J}ac_x\pi(\phi, \bar{\varepsilon})) = \sum_{\bar{\varepsilon}' \in \widehat{\mathcal{S}}_{\phi_-}} \bar{\varepsilon}'(s_-) f_{G_-}(\pi(\phi_-, \bar{\varepsilon}')).$$

In fact (7.10) also holds when $(\rho, 2x+1) \in \text{Jord}(\phi_{II})$, and the argument is similar.

By the linear independence of characters of irreducible smooth representations of $G(F)$, $\pi(\phi_-, \bar{\varepsilon}')$ is in $\bar{J}ac_x\pi(\phi, \bar{\varepsilon})$ only when

$$\bar{\varepsilon}(s) = \bar{\varepsilon}'(s_-)$$

for all $s \in \mathcal{S}_{\phi}$, i.e., $\bar{\varepsilon}' = \bar{\varepsilon}$. This implies $\bar{J}ac_x\pi(\phi, \bar{\varepsilon}) = 0$ for $\bar{\varepsilon} \notin \widehat{\mathcal{S}}_{\phi_-}$. Then after a little thought, one can see

$$\pi(\phi_-, \bar{\varepsilon}) = \bar{J}ac_x\pi(\phi, \bar{\varepsilon})$$

for all $\bar{\varepsilon} \in \widehat{\mathcal{S}}_{\phi_-}$. □

Now we are in the position to prove Theorem 3.3. For the convenience of readers we will restate the theorem here.

Theorem 7.4 (Mœglin). *The Σ_0 -orbits of irreducible supercuspidal representations of $G(F)$ can be parametrized by $\phi \in \bar{\Phi}_2(G)$ and $\bar{\varepsilon} \in \mathcal{S}_{\phi}$ satisfying the following properties:*

- (1) if $(\rho, a) \in \text{Jord}(\phi)$, then $(\rho, a-2) \in \text{Jord}(\phi)$ as long as $a-2 > 0$;
- (2) if $(\rho, a), (\rho, a-2) \in \text{Jord}(\phi)$, then $\varepsilon(\rho, a)\varepsilon(\rho, a-2) = -1$;
- (3) if $(\rho, 2) \in \text{Jord}(\phi)$, then $\varepsilon(\rho, 2) = -1$.

Proof. Let π be an irreducible discrete series representation of $G(F)$, and we can assume $[\pi] = \pi(\phi, \bar{\varepsilon})$ for some $\phi \in \bar{\Phi}_2(G)$ and $\bar{\varepsilon} \in \bar{\mathcal{S}}_\phi$. It is not hard to see that π is supercuspidal if and only if $\bar{\text{Jac}}_x \pi(\phi, \bar{\varepsilon}) = 0$ for any unitary irreducible supercuspidal representation ρ of $GL(d_\rho, F)$ and any real number x . Then by (7.3), it is enough to consider the cases when $(\rho, 2x+1) \in \text{Jord}(\phi)$. Note each of the conditions in this theorem excludes exactly one situation in Lemma 7.3 for $\bar{\text{Jac}}_x \pi(\phi, \bar{\varepsilon}) \neq 0$. So it is clear that these conditions are both necessary and sufficient for π being supercuspidal. $\square$

Remark 7.5. The necessity of condition (1) has already been established by Proposition 3.1, but in this proof we do not need to know that result.

8. CUSPIDAL SUPPORT OF DISCRETE SERIES

In this section we are going to characterize the cuspidal supports of discrete series representations of $G(F)$. Let $\phi \in \bar{\Phi}_2(G)$, for any $(\rho, a) \in \text{Jord}(\phi)$, we denote by a_- the biggest positive integer smaller than a in $\text{Jord}_\rho(\phi)$. And we would also like to write $a_{\min}$ for the minimum of $\text{Jord}_\rho(\phi)$. If $a = a_{\min}$, we let $a_- = 0$ if a is even, and -1 otherwise. In this case, we always assume $\varepsilon(\rho, a)\varepsilon(\rho, a_-) = -1$.

Proposition 8.1. *Suppose $\phi \in \bar{\Phi}_2(G)$, and $\varepsilon \in \widehat{\mathcal{S}}_\phi^{\Sigma_0}$.*

(1) *If $\varepsilon(\rho, a)\varepsilon(\rho, a_-) = -1$ and $a_- < a - 2$, then*

$$(8.1) \quad \pi(\phi, \bar{\varepsilon}) \hookrightarrow \langle (a-1)/2, \dots, (a_-+3)/2 \rangle \rtimes \pi(\phi', \bar{\varepsilon}')$$

is the unique irreducible element in $\bar{\text{Rep}}(G)$ as an $\bar{\mathcal{H}}(G)$ -submodule, where

$$\text{Jord}(\phi') = \text{Jord}(\phi) \cup \{(\rho, a_- + 2)\} \setminus \{(\rho, a)\},$$

and

$$\varepsilon'(\cdot) = \varepsilon(\cdot) \text{ over } \text{Jord}(\phi) \setminus \{(\rho, a)\}, \quad \varepsilon'(\rho, a_- + 2) = \varepsilon(\rho, a).$$

(2) *If $\varepsilon(\rho, a)\varepsilon(\rho, a_-) = 1$, then*

$$(8.2) \quad \pi(\phi, \bar{\varepsilon}) \hookrightarrow \langle (a-1)/2, \dots, -(a_- - 1)/2 \rangle \rtimes \pi(\phi', \bar{\varepsilon}'),$$

where

$$\text{Jord}(\phi') = \text{Jord}(\phi) \setminus \{(\rho, a), (\rho, a_-)\},$$

and $\varepsilon'(\cdot)$ is the restriction of $\varepsilon(\cdot)$. In particular, suppose $\varepsilon_1 \in \widehat{\mathcal{S}}_\phi^{\Sigma_0}$ satisfying $\varepsilon_1(\cdot) = \varepsilon(\cdot)$ over $\text{Jord}(\phi')$ and

$$\varepsilon_1(\rho, a) = -\varepsilon(\rho, a), \quad \varepsilon_1(\rho, a_-) = -\varepsilon(\rho, a_-).$$

If $\bar{\varepsilon}_1 = \bar{\varepsilon}$, then the induced $\bar{\mathcal{H}}(G)$ -module in (8.2) has a unique irreducible element in $\bar{\text{Rep}}(G)$ as an $\bar{\mathcal{H}}(G)$ -submodule. Otherwise, it has two irreducible elements in $\bar{\text{Rep}}(G)$ as $\bar{\mathcal{H}}(G)$ -submodules, namely

$$\pi(\phi, \bar{\varepsilon}) \oplus \pi(\phi, \bar{\varepsilon}_1).$$

(3) *If $\varepsilon(\rho, a_{\min}) = 1$ and $a_{\min}$ is even, then*

$$(8.3) \quad \pi(\phi, \bar{\varepsilon}) \hookrightarrow \langle (a_{\min} - 1)/2, \dots, 1/2 \rangle \rtimes \pi(\phi', \bar{\varepsilon}')$$

is the unique irreducible element in $\bar{\text{Rep}}(G)$ as an $\bar{\mathcal{H}}(G)$ -submodule, where

$$\text{Jord}(\phi') = \text{Jord}(\phi) \setminus \{(\rho, a_{\min})\},$$

and $\varepsilon'(\cdot)$ is the restriction of $\varepsilon(\cdot)$.

Proof. The proofs of part (1) and part (3) are almost the same, so here we will only give the proof of part (1). We start by considering the Jacquet module $\bar{\text{Jac}}_{(a-1)/2, \dots, (a_-+3)/2} \pi(\phi, \bar{\varepsilon})$, and by applying Lemma 7.3 multiple times we have

$$\bar{\text{Jac}}_{(a-1)/2, \dots, (a_-+3)/2} \pi(\phi, \bar{\varepsilon}) = \pi(\phi', \bar{\varepsilon}').$$

It follows from Corollary 5.4 that

$$\pi(\phi, \bar{\varepsilon}) \hookrightarrow \rho^{\frac{a-1}{2}} \times \dots \times \rho^{\frac{a_-+3}{2}} \rtimes \pi(\phi', \bar{\varepsilon}').$$

By Lemma 5.2, we can take an irreducible constituent τ in $\rho||^{\frac{a-1}{2}} \times \cdots \times \rho||^{\frac{a-+3}{2}}$, such that

$$\pi(\phi, \bar{\varepsilon}) \hookrightarrow \tau \rtimes \pi(\phi', \bar{\varepsilon}').$$

So it is enough to show $\tau = < \frac{a-1}{2}, \dots, \frac{a-+3}{2} >$. If this is not the case, we know from Lemma 5.7 that $\text{Jac}_x \tau \neq 0$ for some $(a_- + 3)/2 \leq x < (a - 1)/2$. So $\tau \hookrightarrow \rho||^x \times \tau'$ for some irreducible representation τ' , and

$$\pi(\phi, \bar{\varepsilon}) \hookrightarrow \rho||^x \times \tau' \rtimes \pi(\phi', \bar{\varepsilon}').$$

By Frobenius reciprocity, $\bar{\text{Jac}}_x \pi(\phi, \bar{\varepsilon}) \neq 0$. However, $(\rho, 2x + 1) \notin \text{Jord}(\phi)$ under our assumption, so we get a contradiction (see (7.3)).

To see the induced $\bar{\mathcal{H}}(G)$ -module in (8.1) has a unique irreducible element in $\bar{\text{Rep}}(G)$ as an $\bar{\mathcal{H}}(G)$ -submodule, we can compute its Jacquet module under $\bar{\text{Jac}}_{(a-1)/2, \dots, (a-+3)/2}$. By applying the formula (5.4), we find the Jacquet module consists of

$$\text{Jac}_{X_1} < (a - 1)/2, \dots, (a_- + 3)/2 > \rtimes \text{Jac}_{-X_2}^{op} < (a - 1)/2, \dots, (a_- + 3)/2 > \rtimes \bar{\text{Jac}}_{X_3} \pi(\phi', \bar{\varepsilon}'),$$

where

$$\{(a - 1)/2, \dots, (a_- + 3)/2\} = X_1 \sqcup X_2 \sqcup X_3,$$

and X_i inherits the order from $\{(a - 1)/2, \dots, (a_- + 3)/2\}$. Note $\text{Jac}_{X_1} < (a - 1)/2, \dots, (a_- + 3)/2 > \neq 0$ only if X_1 is a segment $\{(a - 1)/2, \dots, x_1\}$. Similarly, $\text{Jac}_{-X_2}^{op} < (a - 1)/2, \dots, (a_- + 3)/2 > \neq 0$ only if X_2 is a segment $\{-(a_- + 3)/2, \dots, x_2\}$. Since $-(a_- + 3)/2 \notin \{(a - 1)/2, \dots, (a_- + 3)/2\}$, X_2 has to be empty. Therefore the Jacquet module can only contain terms like

$$\text{Jac}_{(a-1)/2, \dots, x_1} < (a - 1)/2, \dots, (a_- + 3)/2 > \rtimes \bar{\text{Jac}}_{x_1-1, \dots, (a-+3)/2} \pi(\phi', \bar{\varepsilon}').$$

But from our definition of $\text{Jord}(\phi')$, we see $\{a, \dots, (a_- + 4)\}$ has no intersection with $\text{Jord}_\rho(\phi')$, so

$$\bar{\text{Jac}}_{x_1-1, \dots, (a-+3)/2} \pi(\phi', \bar{\varepsilon}') = 0$$

by (7.3). Hence we can only have

$$\bar{\text{Jac}}_{(a-1)/2, \dots, (a-+3)/2} \left(< (a - 1)/2, \dots, (a_- + 3)/2 > \rtimes \pi(\phi', \bar{\varepsilon}') \right) = \pi(\phi', \bar{\varepsilon}').$$

Note this implies $< (a - 1)/2, \dots, (a_- + 3)/2 > \rtimes \pi(\phi', \bar{\varepsilon}')$ has a unique irreducible element in $\bar{\text{Rep}}(G)$ as an $\bar{\mathcal{H}}(G)$ -submodule.

For part (2), we will first consider $\bar{\text{Jac}}_{(a-1)/2, \dots, (a-+1)/2} \pi(\phi, \bar{\varepsilon})$, and again by applying Lemma 7.3 multiple times we have

$$\bar{\text{Jac}}_{(a-1)/2, \dots, (a-+1)/2} \pi(\phi, \bar{\varepsilon}) = \pi(\phi_-, \bar{\varepsilon}_-),$$

where $\text{Jord}(\phi_-) = \text{Jord}(\phi) \cup \{(\rho, a_-)\} \setminus \{(\rho, a)\}$, and $\varepsilon_-(\cdot)$ is the restriction of $\varepsilon(\cdot)$ to $\text{Jord}(\phi_-)$ (forgetting multiplicities). As in part (1), we can show from here that

$$(8.4) \quad \pi(\phi, \bar{\varepsilon}) \hookrightarrow < (a - 1)/2, \dots, (a_- + 1)/2 > \rtimes \pi(\phi_-, \bar{\varepsilon}_-).$$

Note $\bar{\Pi}_{\phi_-} = \text{St}(\rho, a_-) \rtimes \bar{\Pi}_{\phi'}$ (see (7.6)), so

$$\pi(\phi_-, \bar{\varepsilon}_-) \hookrightarrow \text{St}(\rho, a_-) \rtimes \pi(\phi', \bar{\varepsilon}') = < (a_- - 1)/2, \dots, -(a_- - 1)/2 > \rtimes \pi(\phi', \bar{\varepsilon}'),$$

and hence

$$(8.5) \quad \pi(\phi, \bar{\varepsilon}) \hookrightarrow < (a - 1)/2, \dots, (a_- + 1)/2 > \rtimes < (a_- - 1)/2, \dots, -(a_- - 1)/2 > \rtimes \pi(\phi', \bar{\varepsilon}').$$

By Lemma 5.2, we can take an irreducible constituent τ in

$$< (a - 1)/2, \dots, (a_- + 1)/2 > \rtimes < (a_- - 1)/2, \dots, -(a_- - 1)/2 >,$$

such that

$$\pi(\phi, \bar{\varepsilon}) \hookrightarrow \tau \rtimes \pi(\phi', \bar{\varepsilon}').$$

Therefore it suffices to show $\tau = \langle (a-1)/2, \dots, -(a_- - 1)/2 \rangle$. If this is not the case, then by Theorem B.1

$$\tau \hookrightarrow \langle (a_- - 1)/2, \dots, -(a_- - 1)/2 \rangle \times \langle (a-1)/2, \dots, (a_- + 1)/2 \rangle.$$

And by Frobenius reciprocity, we have

$$\bar{\text{Jac}}_{(a_- - 1)/2, \dots, -(a_- - 1)/2} \pi \neq 0.$$

But this is impossible, because one can check

$$\text{Jac}_{(a_- - 1)/2, \dots, -(a_- - 1)/2}^\theta \pi \phi = 0.$$

At last, we still need to show the irreducible elements in $\bar{\text{Rep}}(G)$ as submodules of the induced $\bar{\mathcal{H}}(G)$ -module in (8.2) are either $\pi(\phi, \bar{\varepsilon})$ or $\pi(\phi, \bar{\varepsilon}) \oplus \pi(\phi, \bar{\varepsilon}_1)$ depending on whether $\bar{\varepsilon}$ and $\bar{\varepsilon}_1$ are equal or not. Note we can show in the same way as in part (1) that $\pi(\phi, \bar{\varepsilon})$ is the unique irreducible element in $\bar{\text{Rep}}(G)$ as a submodule of the induced $\bar{\mathcal{H}}(G)$ -module in (8.4). And the same is true for $\pi(\phi, \bar{\varepsilon}_1)$. Since $\bar{\varepsilon} = \bar{\varepsilon}_1$ if and only if $\bar{\varepsilon} = \bar{\varepsilon}_{1,-}$, where $\varepsilon_{1,-}(\cdot)$ is again the restriction of $\varepsilon_1(\cdot)$ to $Jord(\phi_-)$ (forgetting multiplicities), let us assume $\bar{\varepsilon} \neq \bar{\varepsilon}_1$ first. Then by (7.7)

$$\pi(\phi_-, \bar{\varepsilon}_-) \oplus \pi(\phi_-, \bar{\varepsilon}_{1,-}) = St(\rho, a_-) \rtimes \pi(\phi', \bar{\varepsilon}'),$$

and hence the irreducible elements in $\bar{\text{Rep}}(G)$ as submodules of the induced $\bar{\mathcal{H}}(G)$ -module in (8.5) are exactly $\pi(\phi, \bar{\varepsilon}) \oplus \pi(\phi, \bar{\varepsilon}_1)$. So we only need to show the induced $\bar{\mathcal{H}}(G)$ -modules in (8.2) and (8.5) have the same irreducible elements in $\bar{\text{Rep}}(G)$ as submodules. One direction is clear, i.e., the irreducible elements in $\bar{\text{Rep}}(G)$ as submodules of

$$\langle (a-1)/2, \dots, (a_- + 1)/2 \rangle \times \langle (a_- - 1)/2, \dots, -(a_- - 1)/2 \rangle \rtimes \pi(\phi', \bar{\varepsilon}')$$

contain that of

$$\langle (a-1)/2, \dots, -(a_- - 1)/2 \rangle \rtimes \pi(\phi', \bar{\varepsilon}')$$

And from what we have shown, it is clear that $\pi(\phi, \bar{\varepsilon}) \oplus \pi(\phi, \bar{\varepsilon}_1)$ are in $\langle (a-1)/2, \dots, -(a_- - 1)/2 \rangle \rtimes \pi(\phi', \bar{\varepsilon}')$, so they have to contain the same irreducible elements in $\bar{\text{Rep}}(G)$ as $\bar{\mathcal{H}}(G)$ -submodules. Now if $\bar{\varepsilon} = \bar{\varepsilon}_1$, we have by (7.7)

$$\pi(\phi_-, \bar{\varepsilon}_-) = St(\rho, a_-) \rtimes \pi(\phi', \bar{\varepsilon}'),$$

and the rest of the argument is the same. □

9. REMARKS ON EVEN ORTHOGONAL GROUPS

The previous results of this paper can also be extended to representations of $G^{\Sigma_0}(F)$. Note the only nontrivial case here is when G is special even orthogonal. First, we will extend Proposition 3.2.

Corollary 9.1. *Suppose π is a supercuspidal representation of $G(F)$ and $[\pi] \in \bar{\Pi}_\phi$ for some $\phi \in \bar{\Phi}_2(G)$. Let π^{Σ_0} be any irreducible representation of $G^{\Sigma_0}(F)$, whose restriction to G contains π . Then for any unitary irreducible supercuspidal representation ρ of $GL(d_\rho, F)$ and real number a_ρ , the parabolic induction*

$$\rho ||^{\pm(a_\rho+1)/2} \rtimes \pi^{\Sigma_0}$$

reduces if and only if ρ is self-dual and

$$(9.1) \quad a_\rho = \begin{cases} \max Jord_\rho(\phi), & \text{if } Jord_\rho(\phi) \neq \emptyset, \\ 0, & \text{if } Jord_\rho(\phi) = \emptyset, \rho \text{ is of opposite type to } \widehat{G}, \\ -1, & \text{otherwise.} \end{cases}$$

Proof. We can assume G is special even orthogonal. First we would like to give the relation of irreducibility between an irreducible representation π of $G(F)$ and an irreducible representation π^{Σ_0} of $G^{\Sigma_0}(F)$ which contains π in its restriction to $G(F)$. For any irreducible representation τ of $GL(d, F)$, it is easy to show the following fact:

- If $\pi \not\cong \pi^{\theta_0}$, $\tau \rtimes \pi^{\Sigma_0}$ is irreducible if and only if $\tau \rtimes \pi$ is irreducible and $(\tau \rtimes \pi)^{\theta_0} \not\cong \tau \rtimes \pi$.

- If $\pi \cong \pi^{\theta_0}$, $\tau \rtimes \pi$ is irreducible if and only if $\tau \rtimes \pi^{\Sigma_0}$ is irreducible and $\tau \rtimes \pi^{\Sigma_0} \not\cong (\tau \rtimes \pi^{\Sigma_0}) \otimes \omega_0$.

Let $\tau = \rho ||^{(a_\rho+1)/2}$ and π be supercuspidal. We assume $\tau \rtimes \pi$ is a representation of $G_+(F)$. Note the condition (3.4) implies (9.1). To see the necessity of the condition (9.1), we need to show if it is not satisfied, then $\tau \rtimes \pi^{\Sigma_0}$ is irreducible. Since $\tau \rtimes \pi$ is irreducible in this case, it suffices to consider $\pi \not\cong \pi^{\theta_0}$, and we would like to show $(\tau \rtimes \pi)^{\theta_0} \not\cong \tau \rtimes \pi$. Since τ and π are both supercuspidal, this is also equivalent to show there does not exist a Weyl group element of $G(F)$ sending $\tau \rtimes \pi^{\theta_0}$ to $\tau \rtimes \pi$, i.e., $\tau \not\cong \tau^\vee$ or d is even. Suppose $\tau \cong \tau^\vee$ and d is odd, then $a_\rho = -1$ and ρ is necessarily of orthogonal type, hence one can only have $Jord_\rho(\phi) \neq \emptyset$ in view of (9.1). This implies $\mathcal{S}_\phi^{\Sigma_0} \neq \mathcal{S}_\phi$, so $\pi \cong \pi^{\theta_0}$ (see the remarks after Theorem 2.3) and we get a contradiction.

To see the reducibility condition (9.1) is also sufficient, we first consider the case $\pi \cong \pi^{\theta_0}$, then the condition (9.1) becomes the same as (3.4). If (9.1) is satisfied, then $\tau \rtimes \pi$ reduces. Suppose $\tau \rtimes \pi^{\Sigma_0}$ is irreducible, then $\tau \rtimes \pi^{\Sigma_0} \cong (\tau \rtimes \pi^{\Sigma_0}) \otimes \omega_0$, and hence

$$(\tau \rtimes \pi^{\Sigma_0})|_{G_+} \cong \tau \rtimes (\pi^{\Sigma_0}|_G) \cong \tau \rtimes \pi \cong \pi_+ \oplus \pi_+^{\theta_0},$$

where $\pi_+ \not\cong \pi_+^{\theta_0}$. By the theory of Langlands quotient, one must have $\tau \cong \rho$. Define ϕ_+ by

$$Jord(\phi_+) := Jord(\phi) \cup \{(\rho, 1) \text{ with multiplicity } 2\}.$$

Then $[\rho \rtimes \pi] \subseteq \bar{\Pi}_{\phi_+}$. Since $\pi \cong \pi^{\theta_0}$, we have $\mathcal{S}_\phi^{\Sigma_0} \neq \mathcal{S}_\phi$, and it follows $\mathcal{S}_{\phi_+}^{\Sigma_0} \neq \mathcal{S}_{\phi_+}$. So $\pi_+^{\theta_0} \cong \pi_+$. This is a contradiction.

At last, we can assume $\pi \not\cong \pi^{\theta_0}$, and it suffices for us to show if $\tau \rtimes \pi^{\Sigma_0}$ is irreducible, then (9.1) is not satisfied. In this case $\tau \rtimes \pi$ is irreducible and $(\tau \rtimes \pi)^{\theta_0} \not\cong \tau \rtimes \pi$. In particular, (3.4) is not satisfied. So we only need to exclude the case that ρ is of orthogonal type, $Jord_\rho(\phi) = \emptyset$, $a_\rho = -1$ and d is odd. In this case $[\tau \rtimes \pi] = [\rho \rtimes \pi] \in \bar{\Pi}_{\phi_+}$ and $\mathcal{S}_{\phi_+}^{\Sigma_0} \neq \mathcal{S}_{\phi_+}$, so $(\tau \rtimes \pi)^{\theta_0} \cong \tau \rtimes \pi$, which again leads to a contradiction. This finishes the proof. $\square$

Next, we would like to extend Lemma 7.3.

Lemma 9.2. Suppose $\phi \in \bar{\Phi}_2(G)$, and $(\rho, 2x+1) \in Jord(\phi)$ with $x > 0$. Let $\phi_- \in \bar{\Phi}_{bdd}(G_-)$ such that

$$Jord(\phi_-) = Jord(\phi) \cup \{(\rho, 2x-1)\} \setminus \{(\rho, 2x+1)\}.$$

Then we have the following facts:

- (1) If $x > 1/2$ and $(\rho, 2x-1) \notin Jord(\phi)$, then $\pi^{\Sigma_0}(\phi_-, \varepsilon) = Jac_x \pi^{\Sigma_0}(\phi, \varepsilon)$ for all $\varepsilon \in \widehat{\mathcal{S}_{\phi_-}^{\Sigma_0}} \cong \widehat{\mathcal{S}_{\phi}^{\Sigma_0}}$.
- (2) If $x > 1/2$ and $(\rho, 2x-1) \in Jord(\phi)$, then $Jac_x \pi^{\Sigma_0}(\phi, \varepsilon) = 0$ unless $\varepsilon \in \widehat{\mathcal{S}_{\phi_-}^{\Sigma_0}}$, i.e.,

$$\varepsilon(\rho, 2x+1)\varepsilon(\rho, 2x-1) = 1,$$

in which case $\pi^{\Sigma_0}(\phi_-, \varepsilon) = Jac_x \pi^{\Sigma_0}(\phi, \varepsilon)$.

- (3) If $x = 1/2$, then $Jac_{1/2} \pi^{\Sigma_0}(\phi, \varepsilon) = 0$ unless $\varepsilon \in \widehat{\mathcal{S}_{\phi_-}^{\Sigma_0}}$, i.e.,

$$\varepsilon(\rho, 2) = 1,$$

in which case $\pi^{\Sigma_0}(\phi_-, \varepsilon) = Jac_x \pi^{\Sigma_0}(\phi, \varepsilon)$.

Proof. Let $G = G(n)$ be special even orthogonal. If $n = d_\rho$, then it suffices to assume $Jord(\phi) = \{(\rho, 2)\}$. In this case, we necessarily have $\varepsilon = 1$ and

$$[(Jac_{1/2} \pi^{\Sigma_0}(\phi, \varepsilon))|_G] = [Jac_{1/2}(\pi^{\Sigma_0}(\phi, \varepsilon)|_G)] = \bar{Jac}_{1/2}(\pi(\phi, \bar{\varepsilon})) = 1.$$

So now we can assume $n \neq d_\rho$. We claim $Jac_x \pi^{\Sigma_0}(\phi, \varepsilon) = 0$ unless $\varepsilon \in \widehat{\mathcal{S}_{\phi_-}^{\Sigma_0}}$, in which case,

$$Jac_x \pi^{\Sigma_0}(\phi, \varepsilon) = \pi^{\Sigma_0}(\phi_-, \varepsilon) \text{ or } \pi^{\Sigma_0}(\phi_-, \varepsilon \varepsilon_0).$$

Suppose $\mathcal{S}_\phi^{\Sigma_0} \neq \mathcal{S}_\phi$, then

$$[(Jac_x \pi^{\Sigma_0}(\phi, \varepsilon))|_G] = [Jac_x(\pi^{\Sigma_0}(\phi, \varepsilon)|_G)] = \bar{Jac}_x \pi(\phi, \bar{\varepsilon}) = 0 \text{ or } \pi(\phi_-, \bar{\varepsilon}),$$

and it is nonzero only when $\varepsilon \in \widehat{\mathcal{S}_{\phi_-}^{\Sigma_0}}$. Suppose $\mathcal{S}_{\phi}^{\Sigma_0} = \mathcal{S}_{\phi}$, then

$$[(\text{Jac}_x \pi^{\Sigma_0}(\phi, \varepsilon))|_G] = [\text{Jac}_x(\pi^{\Sigma_0}(\phi, \varepsilon)|_G)] = 2\bar{\text{Jac}}_x(\pi(\phi, \bar{\varepsilon})) = 0 \text{ or } 2\pi(\phi_-, \bar{\varepsilon}),$$

and it is again nonzero only when $\varepsilon \in \widehat{\mathcal{S}_{\phi_-}^{\Sigma_0}}$. So the claim is clear and it also suffices to show the lemma when $\mathcal{S}_{\phi}^{\Sigma_0} \neq \mathcal{S}_{\phi}$, i.e., $\varepsilon_0 \neq 1$. Let us choose $s^* \in \mathcal{S}_{\phi}^{\Sigma_0}$ such that $\varepsilon_0(s^*) = -1$. Then $s^* \notin \mathcal{S}_{\phi}$. Suppose $(H, \phi_H) \rightarrow (\phi, s^*)$, then we have from (4.5)

$$(9.2) \quad f^H(\phi_H) = \sum_{[\pi] \in \bar{\Pi}_{\phi}} \langle s^*, \pi^{\Sigma_0} \rangle f_G(\pi^{\Sigma_0}) = \sum_{\bar{\varepsilon} \in \widehat{\mathcal{S}_{\phi}}} \varepsilon(s^*) f_G(\pi^{\Sigma_0}(\phi, \varepsilon)),$$

for $f \in C_c^{\infty}(G(F) \rtimes \theta_0)$. In the notation of Section 4, we can write

$$H = G_I \times G_{II} \text{ and } \phi_H = \phi_I \times \phi_{II}$$

Without loss of generality we can assume $(\rho \otimes \eta_{II}, 2x+1) \notin \text{Jord}(\phi_{II})$ and $(\rho \otimes \eta_I, 2x+1) \in \text{Jord}(\phi_I)$. Then by (7.3),

$$\bar{\text{Jac}}_x^{II} \bar{\Pi}_{\phi_{II}} = 0.$$

We let $H_- = H_{I-}$ (see Section 6), and define $\phi_{H_-} = \phi_{I-} \times \phi_{II}$, where

$$\text{Jord}(\phi_{I-}) = \text{Jord}(\phi_I) \cup \{(\rho \otimes \eta_I, 2x-1)\} \setminus \{(\rho \otimes \eta_I, 2x+1)\}.$$

So after applying (6.7) to (9.4), we get

$$(9.3) \quad f^{H_-}(\phi_{H_-}) = \sum_{\bar{\varepsilon} \in \widehat{\mathcal{S}_{\phi_-}}} \varepsilon(s^*) f_{G_-}(\text{Jac}_x \pi^{\Sigma_0}(\phi, \varepsilon)) = \sum_{\bar{\varepsilon} \in \widehat{\mathcal{S}_{\phi_-}}} \varepsilon(s^*) f_{G_-}(\pi^{\Sigma_0}(\phi_-, \varepsilon')),$$

for $f \in C_c^{\infty}(G_-(F) \rtimes \theta_0)$, where $\varepsilon' = \varepsilon$ or $\varepsilon' = \varepsilon \varepsilon_0$. Since $(H_-, \phi_{H_-}) \rightarrow (\phi_-, s_-^*)$, where s_-^* is the image of s^* under the projection $\mathcal{S}_{\phi}^{\Sigma_0} \rightarrow \mathcal{S}_{\phi_-}^{\Sigma_0}$ and $s_-^* \notin \mathcal{S}_{\phi_-}$, we also have

$$f^{H_-}(\phi_{H_-}) = \sum_{\bar{\varepsilon} \in \widehat{\mathcal{S}_{\phi_-}}} \varepsilon(s_-^*) f_{G_-}(\pi^{\Sigma_0}(\phi_-, \varepsilon)),$$

for $f \in C_c^{\infty}(G_-(F) \rtimes \theta_0)$. Combining this identity with (9.3), we get

$$(9.4) \quad \sum_{\bar{\varepsilon} \in \widehat{\mathcal{S}_{\phi_-}}} \varepsilon(s_-^*) f_{G_-}(\pi^{\Sigma_0}(\phi_-, \varepsilon')) = \sum_{\bar{\varepsilon} \in \widehat{\mathcal{S}_{\phi_-}}} \varepsilon(s_-^*) f_{G_-}(\pi^{\Sigma_0}(\phi_-, \varepsilon)).$$

By the linear independence of twisted characters of irreducible smooth representations of $G(F)$, we have

$$\varepsilon(s_-^*) f_{G_-}(\pi^{\Sigma_0}(\phi_-, \varepsilon')) = \varepsilon(s_-^*) f_{G_-}(\pi^{\Sigma_0}(\phi_-, \varepsilon))$$

and hence

$$f_{G_-}(\pi^{\Sigma_0}(\phi_-, \varepsilon')) = f_{G_-}(\pi^{\Sigma_0}(\phi_-, \varepsilon)).$$

This implies $\pi^{\Sigma_0}(\phi_-, \varepsilon') = \pi^{\Sigma_0}(\phi_-, \varepsilon)$, so $\varepsilon = \varepsilon'$.

□

As a consequence of this lemma, we can extend Proposition 8.1. We will follow the same setup in the beginning of Section 8.

Proposition 9.3. *Suppose $\phi \in \bar{\Phi}_2(G)$, and $\varepsilon \in \widehat{\mathcal{S}_{\phi}^{\Sigma_0}}$.*

(1) If $\varepsilon(\rho, a)\varepsilon(\rho, a_-) = -1$ and $a_- < a - 2$, then

$$(9.5) \quad \pi^{\Sigma_0}(\phi, \varepsilon) \hookrightarrow \langle (a-1)/2, \dots, (a_-+3)/2 \rangle \rtimes \pi^{\Sigma_0}(\phi', \varepsilon')$$

as the unique irreducible subrepresentation, where

$$Jord(\phi') = Jord(\phi) \cup \{(\rho, a_- + 2)\} \setminus \{(\rho, a)\},$$

and

$$\varepsilon'(\cdot) = \varepsilon(\cdot) \text{ over } Jord(\phi) \setminus \{(\rho, a)\}, \quad \varepsilon'(\rho, a_- + 2) = \varepsilon(\rho, a).$$

(2) If $\varepsilon(\rho, a)\varepsilon(\rho, a_-) = 1$, then

$$(9.6) \quad \pi^{\Sigma_0}(\phi, \varepsilon) \hookrightarrow \langle (a-1)/2, \dots, -(a_- - 1)/2 \rangle \rtimes \pi^{\Sigma_0}(\phi', \varepsilon'),$$

where

$$Jord(\phi') = Jord(\phi) \setminus \{(\rho, a), (\rho, a_-)\},$$

and $\varepsilon'(\cdot)$ is the restriction of $\varepsilon(\cdot)$. In particular, suppose $\varepsilon_1 \in \widehat{\mathcal{S}}_{\phi}^{\Sigma_0}$ satisfying $\varepsilon_1(\cdot) = \varepsilon(\cdot)$ over $Jord(\phi')$ and

$$\varepsilon_1(\rho, a) = -\varepsilon(\rho, a), \quad \varepsilon_1(\rho, a_-) = -\varepsilon(\rho, a_-).$$

Then the induced representation in (9.6) has two irreducible subrepresentations, namely

$$\pi^{\Sigma_0}(\phi, \varepsilon) \oplus \pi^{\Sigma_0}(\phi, \varepsilon_1).$$

(3) If $\varepsilon(\rho, a_{min}) = 1$ and a_{min} is even, then

$$(9.7) \quad \pi^{\Sigma_0}(\phi, \varepsilon) \hookrightarrow \langle (a_{min} - 1)/2, \dots, 1/2 \rangle \rtimes \pi^{\Sigma_0}(\phi', \varepsilon')$$

as the unique irreducible subrepresentation, where

$$Jord(\phi') = Jord(\phi) \setminus \{(\rho, a_{min})\},$$

and $\varepsilon'(\cdot)$ is the restriction of $\varepsilon(\cdot)$.

The proof of this proposition is almost the same as Proposition 8.1, so we omit it here.

10. CLASSIFICATION OF DISCRETE SERIES

Now we want to characterize the irreducible discrete series representations of $G^{\Sigma_0}(F)$ in terms of their cuspidal supports. For any irreducible discrete series representation $\pi^{\Sigma_0}(\phi, \varepsilon)$ of $G^{\Sigma_0}(F)$, we can associate a triple $(Jord, \pi_{cusp}^{\Sigma_0}, \Delta)$. Here $Jord = Jord(\phi)$ and $\pi_{cusp}^{\Sigma_0}$ is a supercuspidal representation of $G_-^{\Sigma_0}(F)$ which is part of the cuspidal support of π^{Σ_0} . Let us assume $\pi_{cusp}^{\Sigma_0} = \pi^{\Sigma_0}(\phi_{cusp}, \varepsilon_{cusp})$. Finally, Δ is a $\mathbb{Z}_2$ -valued function defined on a subset of

$$Jord \sqcup (Jord \times Jord),$$

i.e., Δ is not defined on $(\rho, a) \in Jord$ with a being odd and $Jord_{\rho}(\phi_{cusp}) \neq \emptyset$; Δ is not defined on pairs $(\rho, a), (\rho', a') \in Jord$ with $\rho \neq \rho'$. Moreover, we require Δ to satisfy the following properties:

- (1) $\Delta(\rho, a)\Delta(\rho, a')^{-1} = \Delta(\rho, a; \rho, a')$,
- (2) $\Delta(\rho, a; \rho, a')\Delta(\rho, a'; \rho, a'') = \Delta(\rho, a; \rho, a'')$,
- (3) $\Delta(\rho, a; \rho, a') = \Delta(\rho, a'; \rho, a)$.

In our case, we can define

$$\Delta(\rho, a) = \varepsilon(\rho, a)$$

for $(\rho, a) \in Jord$ with a being even or $Jord_{\rho}(\phi_{cusp}) = \emptyset$; and

$$\Delta(\rho, a; \rho', a') = \varepsilon(\rho, a)\varepsilon(\rho', a')^{-1}$$

for $(\rho, a), (\rho', a') \in Jord$ with $\rho = \rho'$; otherwise Δ is not defined.

In view of Theorem 3.3 and Proposition 9.3, $(\phi_{cusp}, \varepsilon_{cusp})$ can be constructed from (ϕ, ε) as follows. First we take a maximal sequence of parameters ϕ_i for $1 \leq i \leq k$ such that $\phi_1 = \phi$ and ϕ_{i+1} is obtained from ϕ_i by removing (ρ, a) and (ρ, a_-) , where a_- is the biggest positive integer smaller than a in $Jord_{\rho}(\phi_i)$ and $\varepsilon(\rho, a) = \varepsilon(\rho, a_-)$. Secondly, we remove all $(\rho, a) \in Jord(\phi_k)$, where $a = \min Jord_{\rho}(\phi_k)$ is even and

$\varepsilon(\rho, a) = 1$. We denote the resulting parameter by ϕ_{k+1} and index $Jord_\rho(\phi_{k+1}) = \{a_j\}$ for $j \geq 1$ such that $a_{j+1} > a_j$. Then we can identify $Jord(\phi_{k+1})$ with $Jord(\phi_{cusp})$ by sending (ρ, a_j) to $(\rho, 2j - 1)$ if a_j is odd, or $(\rho, 2j)$ if a_j is even. Let ε_{cusp} be the restriction of ε .

In general, we can consider all triples $(Jord, \pi_{cusp}^{\Sigma_0}, \Delta)$ such that $Jord = Jord(\phi)$ for some $\phi \in \bar{\Phi}_2(G)$, $\pi_{cusp}^{\Sigma_0}$ is some supercuspidal representation of $G_-^{\Sigma_0}(F)$ which is of the same type as $G^{\Sigma_0}(F)$, and Δ satisfies the property that we have mentioned above. Let $Jord_\rho = Jord_\rho(\phi)$. Next we will introduce the concept of **admissibility** for such pairs. Let

$$Jord_\rho^+(\phi_{cusp}) = \begin{cases} Jord_\rho(\phi_{cusp}) \cup \{0\}, & \text{if } a_{\min} = \min Jord_\rho \text{ is even and } \Delta(\rho, a_{\min}) = 1, \\ Jord_\rho(\phi_{cusp}), & \text{otherwise.} \end{cases}$$

Then $(Jord, \pi_{cusp}^{\Sigma_0}, \Delta)$ is called an **admissible triple of alternated type** if

- (1) $\Delta(\rho, a; \rho, a_-) = -1$, if a_- is the biggest positive integer smaller than a in $Jord_\rho$.
- (2) $|Jord_\rho^+(\phi_{cusp})| = |Jord_\rho|$.

We say $(Jord', \pi_{cusp}^{\Sigma_0}, \Delta')$ is subordinated to $(Jord, \pi_{cusp}^{\Sigma_0}, \Delta)$ if $Jord'_\rho = Jord_\rho \setminus \{a, a_-\}$, where $\Delta(\rho, a; \rho, a_-) = 1$, and Δ' is the restriction of Δ . Then $(Jord, \pi_{cusp}^{\Sigma_0}, \Delta)$ is called an **admissible triple** if there exists a sequence of triples $(Jord_i, \pi_{cusp}^{\Sigma_0}, \Delta_i)$ for $1 \leq i \leq k$ such that

- (1) $(Jord, \pi_{cusp}^{\Sigma_0}, \Delta) = (Jord_1, \pi_{cusp}^{\Sigma_0}, \Delta_1)$,
- (2) $(Jord_{i+1}, \pi_{cusp}^{\Sigma_0}, \Delta_{i+1})$ is subordinated to $(Jord_i, \pi_{cusp}^{\Sigma_0}, \Delta_i)$ for $1 \leq i \leq k - 1$,
- (3) $(Jord_k, \pi_{cusp}^{\Sigma_0}, \Delta_k)$ is an admissible triple of alternated type.

Comparing this definition with our construction of $(\phi_{cusp}, \varepsilon_{cusp})$ from (ϕ, ε) , it is easy to see that the triples we associate with irreducible discrete series representations are admissible. On the other hand, from any admissible triple $(Jord, \pi_{cusp}^{\Sigma_0}, \Delta)$ with $Jord = Jord(\phi)$ for some $\phi \in \bar{\Phi}_2(G)$ and $\pi_{cusp}^{\Sigma_0} = \pi^{\Sigma_0}(\phi_{cusp}, \varepsilon_{cusp})$, we can always extend $\varepsilon_{cusp}(\cdot)$ in a unique way to $\varepsilon(\cdot) \in \widehat{\mathcal{S}}_\phi^{\Sigma_0}$ such that the triple is associated with $\pi^{\Sigma_0}(\phi, \varepsilon)$. Therefore we have shown the following theorem.

Theorem 10.1 (Mœglin-Tadić). *There is a one to one correspondence between irreducible discrete series representations of $G^{\Sigma_0}(F)$ and admissible triples $(Jord, \pi_{cusp}^{\Sigma_0}, \Delta)$.*

One can also see how to construct irreducible discrete series representations from admissible triples according to Proposition 9.3. If $(Jord, \pi_{cusp}^{\Sigma_0}, \Delta)$ is an admissible triple of alternated type, let

$$l_\rho : Jord_\rho \longrightarrow Jord_\rho^+(\phi_{cusp})$$

be the monotone bijection. Then the corresponding irreducible discrete series representation π^{Σ_0} can be viewed as the unique irreducible subrepresentation of

$$\left(\prod_{\rho} \left(\prod_{a \in Jord_\rho} < (a-1)/2, \dots, (l_\rho(a)+1)/2 > \right) \right) \rtimes \pi_{cusp}^{\Sigma_0},$$

where the product over $Jord_\rho$ is in the increasing order. If $(Jord, \pi_{cusp}^{\Sigma_0}, \Delta)$ is an admissible triple, we can assume $(Jord', \pi_{cusp}^{\Sigma_0}, \Delta')$ is subordinated to $(Jord, \pi_{cusp}^{\Sigma_0}, \Delta)$, where $Jord'_\rho = Jord_\rho \setminus \{a, a_-\}$. Suppose π'^{Σ_0} corresponds to $(Jord', \pi_{cusp}^{\Sigma_0}, \Delta')$, then

$$< (a-1)/2, \dots, -(a_- - 1)/2 > \rtimes \pi'^{\Sigma_0}$$

has two irreducible subrepresentations, and one will correspond to π^{Σ_0} while the other corresponds to the other extension of Δ' to $Jord$.

11. REMARKS ON THE ORIGINAL APPROACH OF MœGLIN AND TADIĆ

The original approach of Mœglin and Tadić to Theorem 10.1 does not depend on Arthur's theory, i.e., Theorem 2.2, 2.3, 4.2 and 4.4. So the first immediate question becomes how to associate a set of Jordan blocks to every irreducible discrete series representation of $G^{\Sigma_0}(F)$ without assuming Arthur's theory. The answer can be motivated by the following result due to Arthur. It follows from the computation of

the R -group defined by parameters and the fact that they are isomorphic to the representation theoretic R -group (see [Art13, Section 2.4 and Section 6.6]).

Theorem 11.1. *Suppose π^{Σ_0} is an irreducible discrete series representation of $G^{\Sigma_0}(F)$, and $\pi^{\Sigma_0} \in \Pi_{\phi}^{\Sigma_0}$ for some $\phi \in \bar{\Phi}_2(G)$. Then for any self-dual irreducible supercuspidal representation ρ of $GL(d_{\rho}, F)$ and positive integer a , $(\rho, a) \in \text{Jord}(\phi)$ if and only if (ρ, a) is of the same type as $\hat{G}$, and*

$$(11.1) \quad St(\rho, a) \rtimes \pi^{\Sigma_0}$$

is irreducible.

It is clear from this theorem that we can associate every irreducible discrete series representation π^{Σ_0} of $G^{\Sigma_0}(F)$ with a set $\text{Jord}(\pi^{\Sigma_0})$ of Jordan blocks as follows,

$$\text{Jord}(\pi^{\Sigma_0}) := \{(\rho, a) \text{ of the same type as } \hat{G} : \rho \text{ is self-dual supercuspidal, } a \in \mathbb{Z}_{>0} \text{ and (11.1) is irreducible}\}.$$

The next question is about the construction of $\mathbb{Z}_2$ -valued function Δ (see Section 10). In [Mœg02], Mœglin defines Δ over a subset of

$$\text{Jord}(\pi^{\Sigma_0}) \sqcup (\text{Jord}(\pi^{\Sigma_0}) \times \text{Jord}(\pi^{\Sigma_0})),$$

i.e., Δ is not defined on $(\rho, a) \in \text{Jord}(\pi^{\Sigma_0})$ with a being odd and $\text{Jord}_{\rho}(\pi_{\text{cuspidal}}^{\Sigma_0}) \neq \emptyset$; Δ is not defined on pairs $(\rho, a), (\rho', a') \in \text{Jord}(\pi^{\Sigma_0})$ with $\rho \neq \rho'$. Moreover, Δ satisfies those properties that we have described in Section 10. Here we will only mention how to define Δ for pairs $(\rho, a), (\rho, a_-) \in \text{Jord}(\pi^{\Sigma_0})$, where a_- is the biggest positive integer in $\text{Jord}_{\rho}(\pi^{\Sigma_0})$ that is smaller than a , and also for $(\rho, a_{\min}) \in \text{Jord}(\pi^{\Sigma_0})$ with $a_{\min} = \min \text{Jord}_{\rho}(\pi^{\Sigma_0})$ being even. In view of Proposition 9.3, this definition is given in the reversed way, i.e.,

- (1) $\Delta(\rho, a; \rho, a_-) = 1$ if and only if

$$\pi^{\Sigma_0} \hookrightarrow \langle (a-1)/2, \dots, (a_-+1)/2 \rangle \rtimes \pi_-^{\Sigma_0}$$

for some irreducible representation $\pi_-^{\Sigma_0}$ of $G_-^{\Sigma_0}(F)$.

- (2) When $a_{\min}$ is even, $\Delta(\rho, a_{\min}) = 1$ if and only if

$$\pi^{\Sigma_0} \hookrightarrow \langle (a_{\min}-1)/2, \dots, 1/2 \rangle \rtimes \pi_-^{\Sigma_0}$$

for some irreducible representation $\pi_-^{\Sigma_0}$ of $G_-^{\Sigma_0}(F)$.

At last, for $G(n)$ we let $N = 2n + 1$ if G is symplectic, and $N = 2n$ otherwise. Then Mœglin proved the following dimension equality.

Theorem 11.2 (Mœglin [Mœg14]). *Suppose π^{Σ_0} is a discrete series representation of $G^{\Sigma_0}(n, F)$, then*

$$\sum_{(\rho, a) \in \text{Jord}(\pi^{\Sigma_0})} ad_{\rho} = N.$$

This theorem becomes trivial if we know Theorem 2.2 and identify $\text{Jord}(\pi^{\Sigma_0}) = \text{Jord}(\phi)$ under Theorem 11.1. But without assuming all these results of Arthur, this theorem is far from being obvious.

APPENDIX A. LOCAL L-FUNCTION

In this appendix, we give explicit formulas for three different types of local L-functions, i.e., Rankin-Selberg L-function, symmetric square L-function and skew symmetric square L-function. Let F be a p -adic field, and q be the number of elements in the residue field of F .

A.1. Rankin-Selberg L-function. We follow [JPSS83] here. Let π be an irreducible smooth representation of $GL(n, F)$ and σ be an irreducible smooth representation of $GL(m, F)$, the local Rankin-Selberg L-function is denoted by $L(s, \pi \times \sigma)$ for $s \in \mathbb{C}$. It satisfies $L(s, \pi \times \sigma) = L(s, \sigma \times \pi)$ and $L(s, \pi||^t \times \sigma) = L(s + t, \pi \times \sigma)$.

Cuspidal case:

Suppose both π and σ are unitary supercuspidal representations.

- (1) If $n \neq m$, then $L(s, \pi \times \sigma) = 1$;
- (2) If $n = m$, then

$$L(s, \pi \times \sigma) = \prod_t (1 - q^{-(s+it)})^{-1}$$

where the product is over all real numbers t such that $\pi||^{it} \cong \sigma^\vee$.

Discrete series case:

We assume π is $St(\rho, a)$ for an irreducible unitary supercuspidal representations ρ and integer a . Similarly we assume σ is $St(\rho', b)$. If $n \geq m$, then

$$L(s, \pi \times \sigma) = \prod_{i=1}^b L(s + \frac{a+b}{2} - i, \rho \times \rho').$$

Tempered case:

Suppose $\pi = \pi_1 \times \cdots \times \pi_l$ and $\sigma = \sigma_1 \times \cdots \times \sigma_k$, where π_i, σ_j are discrete series representations. Then

$$L(s, \pi \times \sigma) = \prod_{i,j} L(s, \pi_i \times \sigma_j).$$

Non-tempered case:

Let π be the Langlands quotient of the induced representation $\Pi = \pi_1||^{u_1} \times \cdots \times \pi_l||^{u_l}$ for tempered representation π_i and real numbers $u_1 > \cdots > u_l$. Let σ be the Langlands quotient of the induced representation $\Sigma = \sigma_1||^{v_1} \times \cdots \times \sigma_k||^{v_k}$ for tempered representation σ_j and real numbers $v_1 > \cdots > v_k$. Then

$$L(s, \pi \times \sigma) = L(s, \Pi \times \Sigma) = \prod_{i,j} L(s + u_i + v_j, \pi_i \times \sigma_j).$$

A.2. Symmetric square and skew-symmetric square L-functions. We follow [Sha92] here. Let π be an irreducible smooth representation of $GL(n, F)$. The symmetric square (resp. skew-symmetric square) L-function is denoted by $L(s, \pi, S^2)$ (resp. $L(s, \pi, \wedge^2)$). We have $L(s, \pi \times \pi) = L(s, \pi, S^2)L(s, \pi, \wedge^2)$, and $L(s, \pi||^t, R) = L(s + 2t, \pi, R)$ for $R = S^2$ or $\wedge^2$.

Cuspidal case

Suppose π is a unitary supercuspidal representation of $GL(n, F)$.

- (1) $L(s, \pi, \wedge^2) = 1$ unless n is even and some unramified twist of π is self-dual. So let us suppose n is even and π is self-dual. Let S be the set of real numbers t modulo $\frac{\pi}{\ln q}\mathbb{Z}$, such that

$$\int_{Sp_n(F) \backslash GL_n(F)} f({}^t g w^{-1} g w) dg \neq 0,$$

for some $f \in C_c^\infty(GL_n(F))$ defining a matrix coefficient of $\pi||^{it}$. Here ${}^t g$ is the transpose of g and

$$(A.1) \quad w = \begin{pmatrix} 0 & & & 1 \\ & & -1 & \\ & & \ddots & \\ & 1 & & \\ -1 & & & 0 \end{pmatrix}.$$

Then

$$L(s, \pi, \wedge^2) = \prod_{t \in S} (1 - q^{-(s+2it)})^{-1}.$$

- (2) $L(s, \pi, S^2) = 1$ unless some unramified twist of π is self-dual. So let us suppose π is self-dual.
- (a) If n is odd, then

$$L(s, \pi, S^2) = (1 - q^{-rs})^{-1},$$

where r is the maximal integer such that $\pi \cong \pi||^{2\pi i/(r \ln q)}$.

- (b) If n is even,

$$L(s, \pi, S^2) = \prod_{t \in S'} (1 - q^{-(s+2it)})^{-1},$$

where S' is the set of real numbers t modulo $\frac{\pi}{\ln q} \mathbb{Z}$ such that $\pi||^{2it} \cong \pi$ and for any $f \in C_c^\infty(GL_n(F))$ defining a matrix coefficient of $\pi||^{it}$, one has

$$\int_{Sp_n(F) \backslash GL_n(F)} f({}^t g w^{-1} g w) dg = 0.$$

Here w is again given by (A.1) and ${}^t g$ is the transpose of g .

Discrete series case:

We assume π is $St(\rho, a)$ for an irreducible unitary supercuspidal representations ρ and integer a . Set $\pi_i = \rho||^{(a+1)/2-i}$ for $1 \leq i \leq a$.

- (1) Suppose a is even, then

$$\begin{aligned} L(s, \pi, \wedge^2) &= \prod_{i=1}^{a/2} L(s, \pi_i, \wedge^2) L(s, \pi_i||^{-1/2}, S^2), \\ L(s, \pi, S^2) &= \prod_{i=1}^{a/2} L(s, \pi_i, S^2) L(s, \pi_i||^{-1/2}, \wedge^2). \end{aligned}$$

- (2) Suppose a is odd, then

$$\begin{aligned} L(s, \pi, \wedge^2) &= \prod_{i=1}^{(a+1)/2} L(s, \pi_i, \wedge^2) \prod_{i=1}^{(a-1)/2} L(s, \pi_i||^{-1/2}, S^2), \\ L(s, \pi, S^2) &= \prod_{i=1}^{(a+1)/2} L(s, \pi_i, S^2) \prod_{i=1}^{(a-1)/2} L(s, \pi_i||^{-1/2}, \wedge^2). \end{aligned}$$

Tempered case:

Suppose $\pi = \pi_1 \times \cdots \times \pi_l$, where π_i are discrete series representations. Then

$$\begin{aligned} L(s, \pi, \wedge^2) &= \prod_{i=1}^l L(s, \pi_i, \wedge^2) \prod_{1 \leq i < j \leq l} L(s, \pi_i \times \pi_j), \\ L(s, \pi, S^2) &= \prod_{i=1}^l L(s, \pi_i, S^2) \prod_{1 \leq i < j \leq l} L(s, \pi_i \times \pi_j). \end{aligned}$$

Non-tempered case:

Let π be the Langlands quotient of the induced representation $\Pi = \pi_1||^{u_1} \times \cdots \times \pi_l||^{u_l}$ for tempered representation π_i and real numbers $u_1 > \cdots > u_l$. Then

$$\begin{aligned} L(s, \pi, \wedge^2) &= L(s, \Pi, \wedge^2) = \prod_{i=1}^l L(s + 2u_i, \pi_i, \wedge^2) \prod_{1 \leq i < j \leq l} L(s + u_i + u_j, \pi_i \times \pi_j), \\ L(s, \pi, S^2) &= L(s, \Pi, S^2) = \prod_{i=1}^l L(s + 2u_i, \pi_i, S^2) \prod_{1 \leq i < j \leq l} L(s + u_i + u_j, \pi_i \times \pi_j). \end{aligned}$$

APPENDIX B. REDUCIBILITY FOR SOME INDUCED REPRESENTATIONS OF $GL(N)$

We define a **segment** to be a finite length arithmetic progression of real numbers with common difference 1 or -1 , it is completely determined by its endpoints x, y , and hence we denote a segment by $[x, y]$ or $\{x, \dots, y\}$. Let F be a p -adic field and ρ be a unitary irreducible supercuspidal representation of $GL(d_\rho, F)$. The normalized induction

$$\rho||^x \times \dots \times \rho||^y$$

has a unique irreducible subrepresentation, which is denoted by $\langle \rho; x, \dots, y \rangle$ or $\langle x, \dots, y \rangle$. If $x \geq y$, this is called Steinberg representation; if $x < y$, this is called Speh representation.

For any two segments $[x, y]$ and $[x', y']$ such that $(x - y)(x' - y') \geq 0$, we say they are **linked** if as sets $[x, y] \not\subseteq [x', y']$, $[x', y'] \not\subseteq [x, y]$, and $[x, y] \cup [x', y']$ can form a segment after imposing the same order. The following theorem is fundamental in determining the reducibility of an induced representation of $GL(n, F)$.

Theorem B.1 (Zelevinsky [Zel80]). *For unitary irreducible supercuspidal representations ρ, ρ' of general linear groups, and segments $[x, y], [x', y']$ such that $(x - y)(x' - y') \geq 0$,*

$$\langle \rho; x, \dots, y \rangle \times \langle \rho'; x', \dots, y' \rangle$$

is reducible if and only if $\rho \cong \rho'$ and $[x, y], [x', y']$ are linked. In case it is reducible, it consists of the unique irreducible subrepresentations of

$$\langle \rho; x, \dots, y \rangle \times \langle \rho; x', \dots, y' \rangle \quad \text{and} \quad \langle \rho; x', \dots, y' \rangle \times \langle \rho; x, \dots, y \rangle.$$

Remark B.2. In fact, Zelevinsky proved this theorem only when both $x - y \geq 0$ and $x' - y' \geq 0$. Nonetheless, the Aubert involution functor on the Grothendieck group of finite length representations of $GL(n, F)$ will send

$$\langle \rho; x, \dots, y \rangle \times \langle \rho'; x', \dots, y' \rangle \quad \text{to} \quad \langle \rho; y, \dots, x \rangle \times \langle \rho'; y', \dots, x' \rangle$$

up to a sign, and it preserves irreducibility (see [Aub95]). So one can easily extend the original result of Zelevinsky to this theorem.

It is natural to ask for the notion of “link” for two segments $[x, y]$ and $[x', y']$ such that $(x - y)(x' - y') < 0$. To do so, we need to first generalize the notion of “segment”. We define a **generalized segment** to be a matrix

$$\begin{bmatrix} x_{11} & \dots & x_{1n} \\ \vdots & & \vdots \\ x_{m1} & \dots & x_{mn} \end{bmatrix}$$

such that each row is a decreasing (resp. increasing) segment and each column is an increasing (resp. decreasing) segment. The normalized induction

$$\times_{i \in [1, m]} \langle \rho; x_{i1}, \dots, x_{in} \rangle$$

has a unique irreducible subrepresentation, and we denote it by $\langle \rho; \{x_{ij}\}_{m \times n} \rangle$. Moreover,

$$\langle \rho; \{x_{ij}\}_{m \times n} \rangle \cong \langle \rho; \{x_{ij}\}_{m \times n}^T \rangle$$

where $\{x_{ij}\}_{m \times n}^T$ is the transpose of $\{x_{ij}\}_{m \times n}$.

For any two generalized segments $\{x_{ij}\}_{m \times n}$ and $\{y_{ij}\}_{m' \times n'}$ with the same monotone properties for the rows and columns, we say they are **linked** if $[x_{m1}, x_{1n}]$, $[y_{m'1}, y_{1n'}]$ are linked, and the four sides of the rectangle formed by $\{x_{ij}\}_{m \times n}$ do not have inclusive relations with the corresponding four sides of the rectangle formed by $\{y_{ij}\}_{m' \times n'}$ (e.g., $[x_{11}, x_{1n}] \not\subseteq [y_{11}, y_{1n'}]$ and $[x_{11}, x_{1n}] \not\supseteq [y_{11}, y_{1n'}]$, etc). It is easy to check that if $\{x_{ij}\}_{m \times n}$ and $\{y_{ij}\}_{m' \times n'}$ are linked, then $\{x_{ij}\}_{m \times n}^T$ and $\{y_{ij}\}_{m' \times n'}^T$ are also linked. So for generalized segments $\{x_{ij}\}_{m \times n}$ and $\{y_{ij}\}_{m' \times n'}$ with different monotone properties for the rows and columns, we say they are **linked** if $\{x_{ij}\}_{m \times n}^T$ and $\{y_{ij}\}_{m' \times n'}$ are linked, or equivalently $\{x_{ij}\}_{m \times n}$ and $\{y_{ij}\}_{m' \times n'}^T$ are linked. One can check this notion of “link” is equivalent to the one in [MW89].

Example B.3. For any two segments $[x, y]$ and $[x', y']$ such that $(x - y)(x' - y') < 0$, we can view them as generalized segments by taking them as rows, and note they have different monotone properties. So we take

$$[x, y]^T = \begin{bmatrix} x \\ \vdots \\ y \end{bmatrix} \quad \text{and} \quad [x', y'] = [x' \cdots y'].$$

It follows that $[x, y]$ and $[x', y']$ are linked if and only if $[y, x], [x', y']$ are linked, and $x, y \notin [x', y']$ and $x', y' \notin [x, y]$.

The next theorem generalizes Theorem B.1 to the case of generalized segments.

Theorem B.4 (Mœglin-Waldspurger [MW89]). *For unitary irreducible supercuspidal representations ρ, ρ' of general linear groups, and generalized segments $\{x_{ij}\}_{m \times n}, \{y_{ij}\}_{m' \times n'}$,*

$$< \rho; \{x_{ij}\}_{m \times n} > \times < \rho'; \{y_{ij}\}_{m' \times n'} >$$

is irreducible unless $\rho \cong \rho'$ and $\{x_{ij}\}_{m \times n}, \{y_{ij}\}_{m' \times n'}$ are linked.

Let a, b be integers, we define $Sp(St(\rho, a), b)$ to be the unique irreducible subrepresentation of

$$St(\rho, a) ||^{-(b-1)/2} \times St(\rho, a) ||^{-(b-3)/2} \times \cdots \times St(\rho, a) ||^{(b-1)/2}.$$

By the definition one can see $Sp(St(\rho, a), b)$ is given by the following generalized segment

$$\begin{bmatrix} (a-b)/2 & \cdots & 1 - (a+b)/2 \\ \vdots & & \vdots \\ (a+b)/2 - 1 & \cdots & -(a-b)/2 \end{bmatrix}$$

The following result is a reinterpretation of Theorem B.4.

Corollary B.5. *For unitary irreducible supercuspidal representations ρ, ρ' of general linear groups, and integers a, b, a', b' , and real number s ,*

$$Sp(St(\rho, a), b) ||^s \times Sp(St(\rho, a'), b')$$

is irreducible unless $\rho \cong \rho'$, $(a + b + a' + b')/2 + s$ is an integer and

$$|(a - a')/2| + |(b - b')/2| < |s| \leq |(a + a' + b + b')/2| - 1.$$

APPENDIX C. CASSELMAN'S FORMULA AND APPLICATION

Let F be a p -adic field, and G be a quasisplit connected reductive group over F . Let θ be an F -automorphism of G preserving an F -splitting, and we assume θ has order l . Suppose π is an irreducible smooth representation of $G(F)$ such that $\pi \cong \pi^\theta$. Let $A_\pi(\theta)$ be an intertwining operator between π and π^θ , then we can define the twisted character of π to be

$$f_{G^\theta}(\pi) := \text{trace} \int_{G(F)} f(g) \pi(g) dg \circ A_\pi(\theta)$$

for $f \in C_c^\infty(G(F))$. It follows from results of Harish-Chandra [HC99] in the non-twisted case and Clozel [Clo87] in the twisted case, that there exists a locally integrable function $\Theta_\pi^{G^\theta}$ on $G(F)$ such that

$$f_{G^\theta}(\pi) = \int_{G(F)} f(g) \Theta_\pi^{G^\theta}(g) dg.$$

We also call this function the twisted character of π . If $P = MN$ is a θ -stable parabolic subgroup of G , we denote by π_N the unnormalized Jacquet module of π with respect to P , then $\text{Jac}_P \pi = \pi_N \otimes \delta_P^{-1/2}$, where δ_P is the usual modulus character. Note $\pi_N \cong \pi_N^\theta$ and $A_\pi(\theta)$ induces an intertwining operator on π_N .

We would like to extend Jac_P to the space of twisted invariant distributions on $G(F)$. Let $R(G^\theta)$ be the space of finite linear combinations of twisted characters of $G(F)$ and

$$R^*(G^\theta) = \text{Hom}(R(G^\theta), \mathbb{C}).$$

For any $\tau \in R(G^\theta)$ and $f \in C_c^\infty(G(F))$, we define $f_{G^\theta}(\tau)$ by linearity. So we can get a homomorphism

$$C_c^\infty(G(F)) \longrightarrow R^*(G^\theta)$$

by sending f to $l_f(\tau) := f_{G^\theta}(\tau)$. Let us denote the image of this homomorphism by $F_{tr}(G^\theta)$. Since $C_c^\infty(G(F))$ is equipped with a direct limit topology of finite dimensional subspaces, any linear functional on this space is continuous. Moreover, we know the twisted invariant linear functionals on $C_c^\infty(G(F))$ are supported on twisted characters (see [Kaz86, Appendix, Theorem 1] and [Wal17, Section 5.5]), so we can identify the space $\widehat{I}(G^\theta)$ of twisted invariant distributions on $G(F)$ with $\widehat{F}_{tr}(G^\theta) := \text{Hom}(F_{tr}(G^\theta), \mathbb{C})$. Under this identification, we have an inclusion of $R(G^\theta)$ in $\widehat{F}_{tr}(G^\theta)$, which sends τ to $L_\tau(l_f) := l_f(\tau)$ for any $f \in C_c^\infty(G(F))$.

The Jacquet functor induces a homomorphism

$$\text{Jac}_P : R(G^\theta) \longrightarrow R(M^\theta),$$

whose dual is

$$\text{Jac}_P^* : R^*(M^\theta) \longrightarrow R^*(G^\theta)$$

(see [Rog88, Lemma 4.1]). The key step in extending Jac_P to the space of twisted invariant distributions is the following lemma.

Lemma C.1. $\text{Jac}_P^*(F_{tr}(M^\theta)) \subseteq F_{tr}(G^\theta)$.

This lemma was first proved in the nontwisted case (see [BDK86, Proposition 3.2]) and was later extended to the twisted case (see [Rog88, Proposition 7.1]). As a consequence, we have a homomorphism

$$\text{Jac}_P^* : F_{tr}(M^\theta) \longrightarrow F_{tr}(G^\theta),$$

whose dual is

$$\text{Jac}_P : \widehat{F}_{tr}(G^\theta) \longrightarrow \widehat{F}_{tr}(M^\theta).$$

By our previous identifications, this gives

$$(C.1) \quad \text{Jac}_P : \widehat{I}(G^\theta) \longrightarrow \widehat{I}(M^\theta).$$

To see this is really an extension of Jac_P on $R(G^\theta)$, we have the following proposition.

Proposition C.2. *The following diagram commutes.*

$$\begin{array}{ccc} R(G^\theta) & \xrightarrow{\text{Jac}_P} & R(M^\theta) \\ \downarrow & & \downarrow \\ \widehat{I}(G^\theta) & \xrightarrow{\text{Jac}_P} & \widehat{I}(M^\theta). \end{array}$$

Proof. Let $\tau \in R(G^\theta)$ and $h \in C_c^\infty(M(F))$. By definition, we have

$$\text{Jac}_P(L_\tau)(l_h) = L_\tau(\text{Jac}_P^* l_h) = (\text{Jac}_P^* l_h)(\tau) = l_h(\text{Jac}_P \tau) = L_{(\text{Jac}_P \tau)}(l_h).$$

This finishes the proof. □

Let $R(G)^{st}$ be the space of stable finite linear combinations of characters of $G(F)$ and

$$R^*(G)^{st} = \text{Hom}(R(G)^{st}, \mathbb{C}).$$

In the same way, we have

$$(C.2) \quad C_c^\infty(G(F)) \longrightarrow R^*(G)^{st}.$$

Let us denote the image by $F_{tr}(G)^{st}$. The space $\widehat{SI}(G)$ of stable invariant distributions on $G(F)$ are linear functionals of the space of stable orbital integrals of $C_c^\infty(G(F))$. It follows from [Art96, Theorem 6.1 and Theorem 6.2] that one can characterize the space of stable orbital integrals as space of functions on certain subset of $R(G)^{st}$ through (C.2). In particular, the stable invariant distributions are also supported on characters. So we can identify $\widehat{SI}(G)$ with $\widehat{F}_{tr}(G)^{st} := \text{Hom}(F_{tr}(G)^{st}, \mathbb{C})$. By [Hir04, Lemma 2.3], the Jacquet functor induces a homomorphism

$$\text{Jac}_P : R(G)^{st} \longrightarrow R(M)^{st},$$

whose dual is

$$\text{Jac}_P^* : R^*(M)^{st} \longrightarrow R^*(G)^{st}.$$

The following lemma is an analogue of Lemma C.1.

Lemma C.3. $\text{Jac}_P^*(F_{tr}(M)^{st}) \subseteq F_{tr}(G)^{st}$.

Proof. By Lemma C.1, we have a commutative diagram

$$\begin{array}{ccc} F_{tr}(M) & \xrightarrow{\text{Jac}_P^*} & F_{tr}(G) \\ \downarrow & & \downarrow \\ R^*(M)^{st} & \xrightarrow{\text{Jac}_P^*} & R^*(G)^{st}. \end{array}$$

Then this lemma follows from the fact that $F_{tr}(M)^{st}$ and $F_{tr}(G)^{st}$ are images of $F_{tr}(M)$ and $F_{tr}(G)$ respectively. □

As a consequence, we have a homomorphism

$$\text{Jac}_P^* : F_{tr}(M)^{st} \longrightarrow F_{tr}(G)^{st},$$

whose dual is

$$\text{Jac}_P : \widehat{F}_{tr}(G)^{st} \longrightarrow \widehat{F}_{tr}(M)^{st}.$$

By our previous identifications, this gives

$$(C.3) \quad \text{Jac}_P : \widehat{SI}(G) \longrightarrow \widehat{SI}(M).$$

The following proposition is a direct consequence of Proposition C.2.

Proposition C.4. *The following diagram commutes.*

$$\begin{array}{ccc} R(G)^{st} & \xrightarrow{\text{Jac}_P} & R(M)^{st} \\ \downarrow & & \downarrow \\ \widehat{SI}(G) & \xrightarrow{\text{Jac}_P} & \widehat{SI}(M). \end{array}$$

For a strongly θ -regular θ -semisimple element g in $G(F)$, let $h = \prod_{i=1}^l \theta^i(g)$, and one can associate it with a θ -stable parabolic subgroup $P_h = M_h N_h$ by the construction in [Cas77], and $g \in M_h(F)$. Now we can state the twisted version of Casselman's formula.

Theorem C.5. *Suppose π is an irreducible smooth representation of $G(F)$ such that $\pi \cong \pi^\theta$, and g is a strongly θ -regular θ -semisimple element in $G(F)$. Let $h = \prod_{i=1}^l \theta^i(g)$. Then*

$$\Theta_\pi^{G^\theta}(g) = \Theta_{\pi_{N_h}}^{M_h^\theta}(g).$$

Casselman proved this theorem only for $\theta = id$ in [Cas77, Theorem 5.2]. It was generalized to the twisted case by Rogawski in [Rog88, Proposition 7.4]. What we are going to use is the following corollary of this theorem.

Corollary C.6. *Let $P = MN$ be a θ -stable parabolic subgroup of G . Suppose π is an irreducible smooth representation of $G(F)$ such that $\pi \cong \pi^\theta$, and m is a strongly θ -regular θ -semisimple element in $G(F)$, which is also contained in $M(F)$. Then one can choose $z_M \in A_M(F)$ (A_M is the maximal split component of the centre of M) with $|\alpha(z_M)|$ sufficiently small (depending on m) for all roots α in N , such that*

$$\Theta_\pi^{G^\theta}(z_M m) = \Theta_{\pi_N}^{M^\theta}(z_M m).$$

Proof. Let $g = z_M m$, and $h = \prod_{i=1}^l \theta^i(g)$. It is not hard to check from the definition of P_h that $P \supseteq P_h$. Let $P_h^M = M \cap P_h$, it is the parabolic subgroup of M associated with h , and it has Levi component M_h . Let $N_h^M = M \cap N_h$. Then

$$\Theta_\pi^{G^\theta}(g) = \Theta_{\pi_{N_h}}^{M_h^\theta}(g) = \Theta_{(\pi_N)_{N_h^M}}^{M_h^\theta}(g) = \Theta_{\pi_N}^{M^\theta}(g).$$

This finishes the proof. $\square$

As an application of this corollary, we are going to establish diagrams (6.1) and (6.5). First, let us recall the general setup of these diagrams. Let H be a twisted endoscopic group of G , and we assume there is an embedding

$$\xi : {}^L H \rightarrow {}^L G,$$

and $\xi({}^L H) \subseteq \text{Cent}(s, {}^L G)$ and $\widehat{H} \cong \text{Cent}(s, \widehat{G})^0$ for some semisimple $s \in \widehat{G} \rtimes \widehat{\theta}$. We fix $(\widehat{\theta}$ -stable) Γ_F -splittings $(\mathcal{B}_H, \mathcal{T}_H, \{\mathcal{X}_{\alpha_H}\})$ and $(\mathcal{B}_G, \mathcal{T}_G, \{\mathcal{X}_\alpha\})$ for $\widehat{H}$ and $\widehat{G}$ respectively. By taking certain $\widehat{G}$ -conjugate of ξ , we can assume $s \in \mathcal{T}_G \rtimes \widehat{\theta}$ and $\xi(\mathcal{T}_H) = (\mathcal{T}_G^{\widehat{\theta}})^0$ and $\xi(\mathcal{B}_H) \subseteq \mathcal{B}_G$. Let $W_H = W(\widehat{H}, \mathcal{T}_H)$ and $W_{G^\theta} = W(\widehat{G}, \mathcal{T}_G)^{\widehat{\theta}}$, then W_H can be viewed as a subgroup of W_{G^θ} . We also view ${}^L H$ as a subgroup of ${}^L G$ through ξ .

We fix a standard θ -stable parabolic subgroup $P = MN$ of G with standard embedding ${}^L P \hookrightarrow {}^L G$. Then there exists a torus $S \subseteq (\mathcal{T}_G^{\widehat{\theta}})^0$ such that ${}^L M = \text{Cent}(S, {}^L G)$. Let $W_{M^\theta} = W(\widehat{M}, \mathcal{T}_G)^{\widehat{\theta}}$. We define

$$W_{G^\theta}(H, M) := \{w \in W_{G^\theta} \mid \text{Cent}(w(S), {}^L H) \rightarrow W_F \text{ surjective} \}.$$

For any $w \in W_{G^\theta}(H, M)$, let us take $g \in \widehat{G}$ such that $\text{Int}(g)$ induces w . Since $\text{Cent}(w(S), {}^L H) \rightarrow W_F$ is surjective, $g^L P g^{-1} \cap {}^L H$ defines a parabolic subgroup of ${}^L H$ with Levi component $g^L M g^{-1} \cap {}^L H$. So we can choose a standard parabolic subgroup $P'_w = M'_w N'_w$ of H with standard embedding ${}^L P'_w \hookrightarrow {}^L H$ such that ${}^L P'_w$ (resp. ${}^L M'_w$) is $\widehat{H}$ -conjugate to $g^L P g^{-1} \cap {}^L H$ (resp. $g^L M g^{-1} \cap {}^L H$). In particular, M'_w can be viewed as a twisted endoscopic group of M , and the embedding $\xi_{M'_w} : {}^L M'_w \rightarrow {}^L M$ is given by the following diagram:

$$\begin{array}{ccccccc} {}^L P'_w & \longleftarrow & {}^L M'_w & \xrightarrow{\xi_{M'_w}} & {}^L M & \longrightarrow & {}^L P \\ \downarrow & & & & & & \downarrow \\ {}^L H & \xrightarrow{\text{Int}(h)} & {}^L H & \xrightarrow{\xi} & {}^L G & \xleftarrow{\text{Int}(g)} & {}^L G \end{array}$$

where $h \in \widehat{H}$ induces an element in W_H . Note the choice of h is unique up to $\widehat{M}'_w$ -conjugation, and so is $\xi_{M'_w}$. If we change g to $h' g m$, where $h' \in \widehat{H}$ induces an element in W_H and $m \in \widehat{M}$ induces an element in W_{M^θ} , then we still get P'_w , but $\xi_{M'_w}$ changes to $\text{Int}(m^{-1}) \circ \xi_{M'_w}$ up to $\widehat{M}'_w$ -conjugation. To summarize, for any element w in

$$W_H \backslash W_{G^\theta}(H, M) / W_{M^\theta}$$

we can associate a standard parabolic subgroup $P'_w = M'_w N'_w$ of H and a $\widehat{M}$ -conjugacy class of embedding $\xi_{M'_w} : {}^L M'_w \rightarrow {}^L M$.

After this setup, we claim the following diagram commutes.

$$(C.4) \quad \begin{array}{ccc} \widehat{SI}(H) & \longrightarrow & \widehat{I}(G^\theta) \\ \oplus_w \text{Jac}_{P'_w} \downarrow & & \downarrow \text{Jac}_P \\ \bigoplus_w \widehat{SI}(M'_w) & \longrightarrow & \widehat{I}(M^\theta), \end{array}$$

where the sum is over $W_H \backslash W_{G^\theta}(H, M)/W_{M^\theta}$, and the horizontal maps correspond to the twisted spectral endoscopic transfers with respect to ξ on the top and $\xi_{M'_w}$ on the bottom. In fact, it is enough to show the commutativity of this diagram restricting to the subspaces of finite linear combinations of (twisted) characters.

$$(C.5) \quad \begin{array}{ccc} R(H)^{st} & \longrightarrow & R(G^\theta) \\ \oplus_w \text{Jac}_{P'_w} \downarrow & & \downarrow \text{Jac}_P \\ \bigoplus_w R(M'_w)^{st} & \longrightarrow & R(M^\theta), \end{array}$$

where the horizontal maps are given by the restrictions of twisted spectral endoscopic transfers, whose existences are due to [Art96] and [MW16, Appendix].

Lemma C.7. *The commutativity of the diagram (C.5) implies that of (C.4).*

Proof. By taking dual of the diagram (C.5), we get

$$\begin{array}{ccc} R^*(H)^{st} & \longleftarrow & R^*(G^\theta) \\ \oplus_w \text{Jac}_{P'_w}^* \uparrow & & \uparrow \text{Jac}_P^* \\ \bigoplus_w R^*(M'_w)^{st} & \longleftarrow & R^*(M^\theta). \end{array}$$

By restriction, we have

$$\begin{array}{ccc} F_{tr}(H)^{st} & \longleftarrow & F_{tr}(G^\theta) \\ \oplus_w \text{Jac}_{P'_w}^* \uparrow & & \uparrow \text{Jac}_P^* \\ \bigoplus_w F_{tr}(M'_w)^{st} & \longleftarrow & F_{tr}(M^\theta). \end{array}$$

Then the diagram (C.4) is simply its dual. □

As a consequence of this lemma, we only need to show (C.5). To apply Corollary C.6, we need to give another description of the twisted spectral endoscopic transfer. With respect to the embedding ξ , there is a map from the semisimple $H(\bar{F})$ -conjugacy classes of $H(\bar{F})$ to the θ -twisted semisimple $G(\bar{F})$ -conjugacy classes of $G(\bar{F})$ (see [KS99]). If Θ^{G^θ} is a finite linear combination of twisted characters of $G(F)$ and Θ^H is a stable finite linear combination of characters of $H(F)$, then we say Θ^H transfers to Θ^{G^θ} if for any strongly θ -regular θ -semisimple element γ_G in $G(F)$

$$(C.6) \quad \Theta^{G^\theta}(\gamma_G) = \sum_{\gamma_H \rightarrow \gamma_G} \frac{D_H(\gamma_H)^2}{D_{G^\theta}(\gamma_G)^2} \Delta_{G,H}(\gamma_H, \gamma_G) \Theta^H(\gamma_H)$$

where the sum is over $H(\bar{F})$ -conjugacy classes of γ_H in $H(F)$ that map to the θ -twisted $G(\bar{F})$ -conjugacy class of γ_G . In this formula, $\Delta_{G,H}(\cdot, \cdot)$ is the transfer factor (see [KS99]), and it is built into the transfer map introduced in Section 4; $D_H(\cdot)$ and $D_{G^\theta}(\cdot)$ are the (twisted) Weyl discriminants.

From now on let us fix $\gamma_G = \gamma_M$ contained in $M(F)$.

- Lemma C.8.** (1) *If the $H(\bar{F})$ -conjugacy class $\{\gamma_H\}_{H(\bar{F})}$ of γ_H in $H(F)$ maps to the θ -twisted $G(\bar{F})$ -conjugacy class $\{\gamma_G\}_{G(\bar{F})}^\theta$ of γ_G with respect to ξ , then there exists $w \in W_H \backslash W_{G^\theta}(H, M)/W_{M^\theta}$ such that some $H(\bar{F})$ -conjugate $\gamma_{M'_w}$ of γ_H is contained in $M'_w(F)$, and the $M'_w(\bar{F})$ -conjugacy class $\{\gamma_{M'_w}\}_{M'_w(\bar{F})}$ of $\gamma_{M'_w}$ maps to the θ -twisted $M(\bar{F})$ -conjugacy class $\{\gamma_M\}_{M(\bar{F})}^\theta$ of γ_M with respect to $\xi_{M'_w}$.*
- (2) *The correspondence in (1) gives a bijection between $H(\bar{F})$ -conjugacy classes $\{\gamma_H\}_{H(\bar{F})}$ of γ_H in $H(F)$ that map to $\{\gamma_G\}_{G(\bar{F})}^\theta$, and triples $(P'_w, \xi_{M'_w}, \{\gamma_{M'_w}\}_{M'_w(\bar{F})})$ indexed by $w \in W_H \backslash W_{G^\theta}(H, M)/W_{M^\theta}$, where $\{\gamma_{M'_w}\}_{M'_w(\bar{F})}$ maps to $\{\gamma_M\}_{M(\bar{F})}^\theta$.*

Proof. We fix a θ -stable pair (B_M, T_M) in M , where B_M is a Borel subgroup of M defined over $\bar{F}$ and $T_M \subseteq B_M$ is a maximal torus defined over F , such that there exists $\tilde{\gamma}_M = m^{-1}\gamma_M\theta(m) \in T_M$ for $m \in M(\bar{F})$ satisfying $N_\theta(\tilde{\gamma}_M) \in (T_M)_\theta(F)$, where $N_\theta : T_M \rightarrow (T_M)_\theta$. Let $B = B_M N$ and $T = T_M$, then (B, T) is a θ -stable pair in G , and we fix $\tilde{\gamma}_G = \tilde{\gamma}_M$.

Suppose $\{\gamma_H\}_{H(\bar{F})}$ maps to $\{\gamma_G\}_{G(\bar{F})}^\theta$, let $T_H = \text{Cent}(\gamma_H, H)$, then there exists an admissible embedding $\iota : T_H \rightarrow T_\theta$ defined over F , where T_θ is the θ -coinvariant group of T , such that it sends γ_H to $N_\theta(\tilde{\gamma}_G)$. The $\hat{\theta}$ -stable Γ_F -splitting of $\hat{G}$ induces a $\hat{\theta}$ -stable Γ_F -splitting $(\mathcal{B}_M, \mathcal{T}_M, \{\mathcal{X}_{\alpha_M}\})$ of $\hat{M}$. Since $T = T_M$ is contained in M , we can choose isomorphism $(\hat{T}^\theta)^0 \rightarrow (\mathcal{T}_M^\theta)^0 = (\mathcal{T}_G^\theta)^0$ up to W_{M^θ} -conjugation. Then we get an element $w \in W_{G^\theta}$ defined by the following diagram

$$\begin{array}{ccccc} \hat{T}_H & \longrightarrow & (\hat{T}^\theta)^0 & \longrightarrow & (\mathcal{T}_M^\theta)^0 \\ \downarrow & & \downarrow & & \parallel \\ \mathcal{T}_H & \xrightarrow{\xi} & (\mathcal{T}_G^\theta)^0 & \xleftarrow{w} & (\mathcal{T}_G^\theta)^0. \end{array}$$

Recall ${}^L M = \text{Cent}(S, {}^L G)$ for $S \subseteq (\mathcal{T}_G^\theta)^0$, so S is Γ_F -invariant. Since $(\hat{T}^\theta)^0 \rightarrow (\mathcal{T}_M^\theta)^0$ is defined by conjugation in M , the image of S in $(\hat{T}^\theta)^0$ is also Γ_F -invariant. Note $T_H \rightarrow T_\theta$ is defined over F , so the image of S in $\hat{T}_H$ is Γ_F -invariant. It follows $\text{Cent}(w(S), {}^L H) \rightarrow W_F$ is surjective, and hence $w \in W_{G^\theta}(H, M)$. The Γ_F -splitting of $\hat{H}$ induces a Γ_F -splitting $(\mathcal{B}_{M'_w}, \mathcal{T}_{M'_w}, \{\mathcal{X}_{\alpha_{M'_w}}\})$ of $\hat{M}'_w$. Then we have the following diagram

$$\begin{array}{ccc} \mathcal{T}_{M'_w} & \xrightarrow{\xi_{M'_w}} & (\mathcal{T}_M^\theta)^0 \\ \downarrow & & \downarrow w \\ \mathcal{T}_H & \xrightarrow{\xi} & (\mathcal{T}_G^\theta)^0. \end{array}$$

The composition

$$\hat{T}_H \rightarrow \mathcal{T}_H \rightarrow \mathcal{T}_{M'_w}$$

induces a homomorphism $\eta : T_H \rightarrow M'_w$ by H -conjugation, which is determined up to M'_w -conjugation. Since the image S' of S in $\mathcal{T}_{M'_w}$ is Γ_F -invariant and ${}^L M'_w = \text{Cent}(S', {}^L H)$, one can show for any $\sigma \in \Gamma_F$, there exists $m' \in M'_w(\bar{F})$ such that $\sigma(\eta) \circ \eta^{-1} = \text{Int}(m')$. By [Kot82, Corollary 2.2], one can choose η to be defined over F . Let $T_{M'_w} = \eta(T_H)$ and $\gamma_{M'_w} = \eta(\gamma_H)$. It follows from the following commutative diagram

$$\begin{array}{ccccccc} \hat{T}_{M'_w} & \longrightarrow & \mathcal{T}_{M'_w} & \xrightarrow{\xi_{M'_w}} & (\mathcal{T}_M^\theta)^0 & \longrightarrow & (\hat{T}_M^\theta)^0 \\ \downarrow & & \downarrow & & \downarrow w & & \parallel \\ \hat{T}_H & \longrightarrow & \mathcal{T}_H & \xrightarrow{\xi} & (\mathcal{T}_G^\theta)^0 & \longrightarrow & (\hat{T}^\theta)^0 \end{array}$$

that

$$T_{M'_w} \xrightarrow{\eta^{-1}} T_H \xrightarrow{\iota} T_\theta = (T_M)_\theta$$

is an admissible embedding with respect to $\xi_{M'_w}$. So $\{\gamma_{M'_w}\}_{M'_w(\bar{F})}$ maps to $\{\gamma_M\}_{M(\bar{F})}^\theta$. This proves part (1) of the lemma.

To prove part (2), we first want to show the map obtained above by sending $\{\gamma_H\}_{H(\bar{F})}$ to the triple $(P'_w, \xi_{M'_w}, \{\gamma_{M'_w}\}_{M'_w(\bar{F})})$ is well-defined. If we fix the admissible embedding $T_H \rightarrow T_\theta$, but change $\hat{T}_H \rightarrow \mathcal{T}_H$ by W_H -conjugation and $(\hat{T}^\theta)^0 \rightarrow (\mathcal{T}_M^\theta)^0$ by W_{M^θ} -conjugation, one can see w is well-defined in $W_H \backslash W_{G^\theta}(H, M) / W_{M^\theta}$. Moreover, $\{\gamma_{M'_w}\}_{M'_w(\bar{F})}$ is uniquely determined. Now let us vary the admissible embedding ι by choosing $\gamma'_H \in \{\gamma_H\}_{H(\bar{F})} \cap H(F)$ and $\iota' : T'_H \rightarrow T_\theta$ defined over F , such that $\gamma'_H \in T'_H(F)$ is again sent to $N_\theta(\tilde{\gamma}_G)$. Suppose $\gamma'_H = h\gamma_H h^{-1}$ for $h \in H(\bar{F})$. We claim

$$\iota^{-1} \circ \iota' : T'_H \rightarrow T_H$$

can be given by conjugation in H . To see this, we consider the following commutative diagram

$$\begin{array}{ccccc} T_H & \longrightarrow & T'_H & \xrightarrow{\text{Int}(h)^{-1}} & T_H \\ \downarrow \iota & & \downarrow \iota' & & \downarrow \iota \\ T_\theta & \xlongequal{\quad} & T_\theta & \xrightarrow{w'} & T_\theta. \end{array}$$

Here w' can be viewed as in $W(G, T)^\theta$. Since θ preserves a splitting of G , we can choose $g \in G(\bar{F})$ such that $\theta(g) = g$ and $\text{Int}(g) = w'$. Since w' fixes $N_\theta(\tilde{\gamma}_G)$, there exists $t \in T(\bar{F})$ such that

$$g\tilde{\gamma}_G g^{-1} = \tilde{\gamma}_G t^{-1} \theta(t).$$

Then

$$(tg)\tilde{\gamma}_G \theta(tg)^{-1} = \tilde{\gamma}_G.$$

Since γ_G is strongly θ -regular, $tg \in T(\bar{F})$ and hence $g \in T(\bar{F})$. It follows $w' = 1$, and we have $\iota^{-1} \circ \iota' = \text{Int}(h)^{-1}$.

As a consequence of our claim, we have

$$\begin{array}{ccccccc} \hat{T}'_H & \longrightarrow & \hat{T}_H & \longrightarrow & (\hat{T}^\theta)^0 & \longrightarrow & (\mathcal{T}_M^\theta)^0 \\ \downarrow & & \downarrow & & \downarrow & & \parallel \\ \mathcal{T}_H & \xlongequal{\quad} & \mathcal{T}_H & \xrightarrow{\xi} & (\mathcal{T}_G^\theta)^0 & \xleftarrow{w} & (\mathcal{T}_G^\theta)^0 \end{array}$$

This means w is unchanged in $W_H \backslash W_{G^\theta}(H, M) / W_{M^\theta}$. Moreover, we have

$$\begin{array}{ccccccc} \hat{T}'_{M'_w} & \xlongequal{\quad} & \hat{T}_{M'_w} & \longrightarrow & \mathcal{T}_{M'_w} & \xrightarrow{\xi_{M'_w}} & (\mathcal{T}_M^\theta)^0 \longrightarrow (\hat{T}_M^\theta)^0 \\ \downarrow & & \downarrow & & \downarrow & & \downarrow w \\ \hat{T}'_H & \longrightarrow & \hat{T}_H & \longrightarrow & \mathcal{T}_H & \xrightarrow{\xi} & (\mathcal{T}_G^\theta)^0 \longrightarrow (\hat{T}^\theta)^0 \end{array}$$

It is easy to see that one still gets the same $\{\gamma_{M'_w}\}_{M'_w(\bar{F})}$.

Next we want to construct the inverse. For any triple $(P'_w, \xi_{M'_w}, \{\gamma_{M'_w}\}_{M'_w(\bar{F})})$, let $T_{M'_w} = \text{Cent}(\gamma_{M'_w}, M'_w)$. Since $\{\gamma_{M'_w}\}_{M'_w(\bar{F})}$ maps to $\{\gamma_M\}_{M(\bar{F})}^\theta$, we have an admissible embedding $T_{M'_w} \rightarrow (T_M)_\theta$ with respect to

$\xi_{M'_w}$, which sends $\gamma_{M'_w}$ to $\tilde{\gamma}_M$. Let $\gamma_H = \gamma_{M'_w}$ and $T_H = T_{M'_w}$. It follows from the following diagram

$$\begin{array}{ccccccc} \widehat{T}_{M'_w} & \longrightarrow & \mathcal{T}_{M'_w} & \xrightarrow{\xi_{M'_w}} & (\mathcal{T}_M^{\widehat{\theta}})^0 & \longrightarrow & (\widehat{T}_M^{\widehat{\theta}})^0 \\ \parallel & & \downarrow & & \downarrow w & & \parallel \\ \widehat{T}_H & \longrightarrow & \mathcal{T}_H & \xrightarrow{\xi} & (\mathcal{T}_G^{\widehat{\theta}})^0 & \longrightarrow & (\widehat{T}^{\widehat{\theta}})^0 \end{array}$$

that $T_H \rightarrow T_\theta$ is an admissible embedding with respect to ξ . So $\{\gamma_H\}_{H(\bar{F})}$ maps to $\{\gamma_G\}_{G(\bar{F})}^\theta$. In this way, we send $(P'_w, \xi_{M'_w}, \{\gamma_{M'_w}\}_{M'_w(\bar{F})})$ to $\{\gamma_H\}_{H(\bar{F})}$, where $\gamma_H = \gamma_{M'_w}$. Finally, it is easy to check that this map does give the inverse. $\square$

As a consequence of this lemma, we can rewrite the right hand side of (C.6) as

$$\sum_w \sum_{\gamma_{M'_w} \rightarrow \gamma_M} \frac{D_H(\gamma_{M'_w})^2}{D_{G^\theta}(\gamma_M)^2} \Delta_{G,H}(\gamma_{M'_w}, \gamma_M) \Theta^H(\gamma_{M'_w}).$$

So the next step is to write the summands in terms of M and M'_w . First, it is easy to check from the definition of transfer factors that

$$\frac{D_H(\gamma_{M'_w})}{D_{G^\theta}(\gamma_M)} \Delta_{G,H}(\gamma_{M'_w}, \gamma_M) = \frac{D_{M'_w}(\gamma_{M'_w})}{D_{M^\theta}(\gamma_M)} \Delta_{M,M'_w}(\gamma_{M'_w}, \gamma_M)$$

Secondly, there is a natural homomorphism from A_M to $A_{M'_w}$ with respect to $\xi_{M'_w}$ (see [KS99]). For $z_M \in A_M(F)$, let $z_{M'_w}$ be its image in $A_{M'_w}(F)$, then the $M'_w(\bar{F})$ -conjugacy class of $z_{M'_w} \gamma_{M'_w}$ maps to the $M(\bar{F})$ -conjugacy class of $z_M \gamma_M$. Moreover, $\sup|\alpha'(z_{M'_w})|$ for roots α' in N'_w is less than $\sup|\prod_{i=1}^l \theta^i(\alpha)(z_M)|$ for roots α in N .

Let

$$\begin{aligned} \Theta^{M^\theta} &= \sum_i c_i \cdot \Theta_{\pi_i}^{M^\theta} \\ (\text{resp. } \Theta^{M'_w} &= \sum_j d_{w,j} \cdot \Theta_{\pi'_{w,j}}^{M'_w}) \end{aligned}$$

be the (resp. stable) finite linear combination of (twisted) characters of M (resp. M'_w) obtained from Θ^{G^θ} (resp. Θ^H) by taking the unnormalized Jacquet modules. If we take $z_M \in A_M(F)$ so that $\sup|\alpha(z_M)|$ is sufficiently small for all roots α in N , then by Corollary C.6,

$$\Theta^{G^\theta}(z_M \gamma_M) = \Theta^{M^\theta}(z_M \gamma_M),$$

and

$$\Theta^H(z_{M'_w} \gamma_{M'_w}) = \Theta^{M'_w}(z_{M'_w} \gamma_{M'_w}).$$

Besides, it is easy to verify

$$D_{G^\theta}(z_M \gamma_M) = \delta_P(z_M \gamma_M)^{-1/2} \cdot D_{M^\theta}(z_M \gamma_M),$$

and

$$D_H(z_{M'_w} \gamma_{M'_w}) = \delta_{P'}(z_{M'_w} \gamma_{M'_w})^{-1/2} \cdot D_{M'_w}(z_{M'_w} \gamma_{M'_w}).$$

Putting all these together, we get

$$\begin{aligned} & \delta_P(z_M \gamma_M)^{-1/2} \Theta^{M^\theta}(z_M \gamma_M) \\ &= \sum_w \sum_{\gamma_{M'_w} \rightarrow \gamma_M} \frac{D_{M'_w}(z_{M'_w} \gamma_{M'_w})^2}{D_{M^\theta}(z_M \gamma_M)^2} \Delta_{M,M'_w}(z_{M'_w} \gamma_{M'_w}, z_M \gamma_M) \delta_{P'}(z_{M'_w} \gamma_{M'_w})^{-1/2} \Theta^{M'_w}(z_{M'_w} \gamma_{M'_w}). \end{aligned}$$

Since $z_M, z_{M'_w}$ are in the centres of M and M'_w respectively, we have

$$\Delta_{M,M'_w}(z_{M'_w} \gamma_{M'_w}, z_M \gamma_M) = \chi'_w(z_M) \Delta_{M,M'_w}(\gamma_{M'_w}, \gamma_M),$$

$$\begin{aligned}
\delta_P(z_M \gamma_M)^{-1/2} \Theta_{\pi_i}^{M^\theta}(z_M \gamma_M) &= \zeta_{M,i}(z_M) \delta_P(\gamma_M)^{-1/2} \Theta_{\pi_i}^{M^\theta}(\gamma_M), \\
\delta_{P'_w}(z_{M'_w} \gamma_{M'_w})^{-1/2} \Theta_{\pi'_{w,j}}^{M'_w}(z_{M'_w} \gamma_{M'_w}) &= \zeta_{M'_w,j}(z_{M'_w}) \delta_{P'_w}(\gamma_{M'_w})^{-1/2} \Theta_{\pi'_{w,j}}^{M'_w}(\gamma_{M'_w}), \\
D_{M^\theta}(z_M \gamma_M) &= D_{M^\theta}(\gamma_M), \\
D_{M'_w}(z_{M'_w} \gamma_{M'_w}) &= D_{M'_w}(\gamma_{M'_w}),
\end{aligned}$$

where $\chi'_w, \zeta_{M,i}, \zeta_{M'_w,j}$ are the corresponding central characters. Hence

$$\begin{aligned}
&\sum_i (c_i \cdot \delta_P(\gamma_M)^{-1/2} \Theta_{\pi_i}^{M^\theta}(\gamma_M)) \cdot \zeta_{M,i}(z_M) \\
&= \sum_w \sum_j (d_{w,j} \sum_{\gamma_{M'_w} \rightarrow \gamma_M} \frac{D_{M'_w}(\gamma_{M'_w})^2}{D_{M^\theta}(\gamma_M)^2} \Delta_{M,M'_w}(\gamma_{M'_w}, \gamma_M) \delta_{P'_w}(\gamma_{M'_w})^{-1/2} \Theta_{\pi'_{w,j}}^{M'_w}(\gamma_{M'_w})) \cdot \chi'_w(z_M) \zeta_{M'_w,j}(z_{M'_w}).
\end{aligned}$$

Let

$$a_{M,i} = c_i \cdot \delta_P(\gamma_M)^{-1/2} \Theta_{\pi_i}^{M^\theta}(\gamma_M),$$

and

$$b_{w,j} = d_{w,j} \sum_{\gamma_{M'_w} \rightarrow \gamma_M} \frac{D_{M'_w}(\gamma_{M'_w})^2}{D_{M^\theta}(\gamma_M)^2} \Delta_{M,M'_w}(\gamma_{M'_w}, \gamma_M) \delta_{P'_w}(\gamma_{M'_w})^{-1/2} \Theta_{\pi'_{w,j}}^{M'_w}(\gamma_{M'_w}).$$

We also write $\chi_{M,i}(z_M) = \zeta_{M,i}(z_M)$ and $\chi_{w,j}(z_M) = \chi'_w(z_M) \zeta_{M'_w,j}(z_{M'_w})$. Then we can get a short expression

$$(C.7) \quad \sum_i a_{M,i} \cdot \chi_{M,i}(z_M) = \sum_w \sum_j b_{w,j} \cdot \chi_{w,j}(z_M),$$

and it suffices for us to show this holds when $z_M = 1$, i.e.,

$$\sum_i a_{M,i} = \sum_w \sum_j b_{w,j}.$$

In fact, we can choose $z_M \in F^\times \hookrightarrow A_M(F)$ such that (C.7) holds provided $|z_M| < q_F^{-k}$ for some positive integer k , where q_F is the order of the residue field of F . Then it is enough to have the following lemma.

Lemma C.9. *For quasicharacters χ_i of $F^\times$ and complex numbers a_i , if*

$$\sum_{i=1}^r a_i \chi_i(z) = 0$$

provided $|z| < q_F^{-k}$ for some positive integer k , then

$$\sum_{i=1}^r a_i = 0.$$

Proof. Suppose χ_i are distinct, we claim $a_i = 0$ for $1 \leq i \leq r$. It is clear that this lemma will follow from our claim. So next we will show the claim by induction on r . When $r = 1$, there is nothing to show. In general, let us first assume all χ_i are unramified. We choose $z_0 \in F^\times$ such that $|z_0| < q_F^{-k}$, and denote $\chi_i(z_0)$ by C_i . Then

$$\sum_i a_i C_i^j = \sum_i a_i \chi_i(z_0^j) = 0$$

for any positive integer j . In particular, $\{a_i\}$ forms a solution of the linear system of equations defined by the matrix $\{C_i^j\}_{r \times r}^T$, where we let $1 \leq j \leq r$. Since $|\det(\{C_i^j\}_{r \times r})| = |\prod_i C_i| \cdot \prod_{i \leq j} |C_i - C_j| \neq 0$, then a_i have to be all zero. Now suppose some χ_i is ramified, we can replace χ_i by $\chi'_i := \chi_i / \chi_1$ for all i . If χ'_i are all unramified, then we are back to the previous case. If χ'_{i_0} is ramified for some $i_0 > 1$, then we can

choose some unit of the ring of integers of F such that $\chi'_{i_0}(u) \neq 1$. By subtracting $\chi_1(u) \sum_i a_i \chi_i(z)$ from $\sum_i a_i \chi_i(uz)$, we get

$$\sum_{i>1} a_i (\chi_i(u) - \chi_1(u)) \chi_i(z) = 0$$

provided $|z| < q_F^{-k}$. By induction, we have $a_i (\chi_i(u) - \chi_1(u)) = 0$ for $i > 1$. Since $\chi_{i_0}(u) - \chi_1(u) \neq 0$ by our assumption, this implies $a_{i_0} = 0$. Hence

$$\sum_{i \neq i_0} a_i \chi_i(z) = 0$$

provided $|z| < q_F^{-k}$. By induction again, we have $a_i = 0$ for $i \neq i_0$. This finishes the proof of the claim. $\square$

REFERENCES

- [Art96] J. Arthur, *On local character relations*, Selecta Math. (N.S.) **2** (1996), no. 4, 501–579.
- [Art13] ———, *The endoscopic classification of representations: orthogonal and symplectic groups*, Colloquium Publications, vol. 61, American Mathematical Society, 2013.
- [Aub95] A.-M. Aubert, *Dualité dans le groupe de Grothendieck de la catégorie des représentations lisses de longueur finie d'un groupe réductif p -adique*, Trans. Amer. Math. Soc. **347** (1995), no. 6, 2179–2189.
- [Ban99] D. Ban, *Parabolic induction and Jacquet modules of representations of $O(2n, F)$* , Glas. Mat. Ser. III **34**(54) (1999), no. 2, 147–185. MR 1739616
- [BDK86] J. Bernstein, P. Deligne, and D. Kazhdan, *Trace Paley-Wiener theorem for reductive p -adic groups*, J. Analyse Math. **47** (1986), 180–192. MR 874050
- [BW00] A. Borel and N. Wallach, *Continuous cohomology, discrete subgroups, and representations of reductive groups*, second ed., Mathematical Surveys and Monographs, vol. 67, American Mathematical Society, Providence, RI, 2000. MR 1721403
- [BZ77] I. N. Bernstein and A. V. Zelevinsky, *Induced representations of reductive p -adic groups. I*, Ann. Sci. École Norm. Sup. (4) **10** (1977), no. 4, 441–472.
- [Cas77] W. Casselman, *Characters and Jacquet modules*, Math. Ann. **230** (1977), no. 2, 101–105.
- [Clo87] L. Clozel, *Characters of nonconnected, reductive p -adic groups*, Canad. J. Math. **39** (1987), no. 1, 149–167.
- [HC99] Harish-Chandra, *Admissible invariant distributions on reductive p -adic groups*, University Lecture Series, vol. 16, American Mathematical Society, Providence, RI, 1999, Preface and notes by Stephen DeBacker and Paul J. Sally, Jr.
- [Hen00] G. Henniart, *Une preuve simple des conjectures de Langlands pour $GL(n)$ sur un corps p -adique*, Invent. Math. **139** (2000), no. 2, 439–455.
- [Hir04] K. Hiraga, *On functoriality of Zelevinski involutions*, Compositio Math. **140** (2004), 1625–1656.
- [HT01] M. Harris and R. Taylor, *The geometry and cohomology of some simple Shimura varieties*, Annals of Mathematics Studies, vol. 151, Princeton University Press, Princeton, NJ, 2001, With an appendix by Vladimir G. Berkovich.
- [Jan06] C. Jantzen, *Jacquet modules of induced representations for p -adic special orthogonal groups*, J. Algebra **305** (2006), no. 2, 802–819.
- [JPSS83] H. Jacquet, I. I. Piatetskii-Shapiro, and J. A. Shalika, *Rankin-Selberg convolutions*, Amer. J. Math. **105** (1983), no. 2, 367–464.
- [Kaz86] D. Kazhdan, *Cuspidal geometry of p -adic groups*, J. Analyse Math. **47** (1986), 1–36. MR 874042
- [Kot82] R. E. Kottwitz, *Rational conjugacy classes in reductive groups*, Duke Math. J. **49** (1982), no. 4, 785–806.
- [KS99] R. E. Kottwitz and D. Shelstad, *Foundations of twisted endoscopy*, Astérisque **55** (1999), no. 255, vi+190.
- [Mœg02] C. Mœglin, *Sur la classification des séries discrètes des groupes classiques p -adiques: paramètres de Langlands et exhaustivité*, J. Eur. Math. Soc. (JEMS) **4** (2002), no. 2, 143–200.
- [Mœg11] ———, *Multiplicité 1 dans les paquets d'Arthur aux places p -adiques*, On certain L -functions, Clay Math. Proc., vol. 13, Amer. Math. Soc., Providence, RI, 2011, pp. 333–374.
- [Mœg14] ———, *Paquets stables des séries discrètes accessibles par endoscopie tordue; leur paramètre de Langlands*, Automorphic forms and related geometry: assessing the legacy of I. I. Piatetski-Shapiro, Contemp. Math., vol. 614, Amer. Math. Soc., Providence, RI, 2014, pp. 295–336.
- [MT02] C. Mœglin and M. Tadić, *Construction of discrete series for classical p -adic groups*, J. Amer. Math. Soc. **15** (2002), no. 3, 715–786 (electronic).
- [MT11] G. Muić and M. Tadić, *Unramified unitary duals for split classical p -adic groups; the topology and isolated representations*, On certain L -functions, Clay Math. Proc., vol. 13, Amer. Math. Soc., Providence, RI, 2011, pp. 375–438.
- [MW89] C. Mœglin and J.-L. Waldspurger, *Le spectre résiduel de $GL(n)$* , Ann. Sci. École Norm. Sup. (4) **22** (1989), no. 4, 605–674.

- [MW06] ———, *Sur le transfert des traces d'un groupe classique p -adique à un groupe linéaire tordu*, Selecta Math. (N.S.) **12** (2006), no. 3-4, 433–515.
- [MW16] ———, *Stabilisation de la formule des traces tordue*, Progress in Mathematics, vol. 316/317, Birkhäuser Basel, 2016.
- [Ngô10] B. C. Ngô, *Le lemme fondamental pour les algèbres de Lie*, Publ. Math. Inst. Hautes Études Sci. (2010), no. 111, 1–169.
- [Rog88] J. D. Rogawski, *Trace Paley-Wiener theorem in the twisted case*, Trans. Amer. Math. Soc. **309** (1988), no. 1, 215–229. MR 957068
- [Sch13] P. Scholze, *The local Langlands correspondence for GL_n over p -adic fields*, Invent. Math. **192** (2013), no. 3, 663–715.
- [Sha81] F. Shahidi, *On certain L -functions*, Amer. J. Math. **103** (1981), no. 2, 297–355.
- [Sha92] ———, *Twisted endoscopy and reducibility of induced representations for p -adic groups*, Duke Math. J. **66** (1992), no. 1, 1–41.
- [Sil79] A. J. Silberger, *Introduction to harmonic analysis on reductive p -adic groups*, Mathematical Notes, vol. 23, Princeton University Press, Princeton, N.J., 1979, Based on lectures by Harish-Chandra at the Institute for Advanced Study, 1971–1973.
- [Tad86] M. Tadić, *Classification of unitary representations in irreducible representations of general linear group (non-Archimedean case)*, Ann. Sci. École Norm. Sup. (4) **19** (1986), no. 3, 335–382.
- [Tad09] ———, *On reducibility and unitarizability for classical p -adic groups, some general results*, Canad. J. Math. **61** (2009), no. 2, 427–450.
- [Wal95] J.-L. Waldspurger, *Une formule des traces locale pour les algèbres de Lie p -adiques*, J. Reine Angew. Math. **465** (1995), 41–99.
- [Wal97] ———, *Le lemme fondamental implique le transfert*, Compositio Math. **105** (1997), no. 2, 153–236.
- [Wal03] ———, *La formule de Plancherel pour les groupes p -adiques (d'après Harish-Chandra)*, J. Inst. Math. Jussieu **2** (2003), no. 2, 235–333.
- [Wal06] ———, *Endoscopie et changement de caractéristique*, J. Inst. Math. Jussieu **5** (2006), no. 3, 423–525.
- [Wal08] ———, *L'endoscopie tordue n'est pas si tordue*, Mem. Amer. Math. Soc. **194** (2008), no. 908, x+261.
- [Wal17] ———, *La formule des traces locale tordue*, à paraître aux Memoirs of AMS (2017).
- [Zel80] A. V. Zelevinsky, *Induced representations of reductive p -adic groups. II. On irreducible representations of $GL(n)$* , Ann. Sci. École Norm. Sup. (4) **13** (1980), no. 2, 165–210.

DEPARTMENT OF MATHEMATICS AND STATISTICS, UNIVERSITY OF CALGARY, CANADA
E-mail address: bin.xu2@ucalgary.ca